

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 31/55	A1	(11) International Publication Number: WO 95/14470 (43) International Publication Date: 1 June 1995 (01.06.95)
<p>(21) International Application Number: PCT/US94/13364</p> <p>(22) International Filing Date: 18 November 1994 (18.11.94)</p> <p>(30) Priority Data: 156,184 22 November 1993 (22.11.93) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 156,184 (CON) Filed on 22 November 1993 (22.11.93)</p> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SANGUINETTI, Michael, C. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). LYNCH, Joseph, J., Jr. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). SALATA, Joseph, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>		<p>(74) Common Representative: MERCK & CO., INC.; Patent Dept., 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: METHODS OF TREATING CARDIAC ARRHYTHMIA</p> <p>(57) Abstract Benzodiazepine analogs have been found to be useful in treating cardiac abnormalities.</p>		

B13

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

TITLE OF THE INVENTION
METHODS OF TREATING CARDIAC ARRHYTHMIA

CROSS REFERENCE

5 This is a continuation in part of U.S. Patent Application Serial No. 08/018,912 which was filed on February 17, 1993, which is a continuation of U.S. Patent Application Serial No. 07/802,000 which was filed on December 3, 1991.

10 BACKGROUND OF THE INVENTION

Cardiac arrhythmias often occur as complications to cardiac diseases such as myocardial infarction and heart failure. In serious cases, arrhythmias give rise to ventricular fibrillation and can cause sudden death.

15 Although various antiarrhythmic agents are now available on the market, those having both satisfactory efficacy and a high margin of safety have not been obtained. For example, antiarrhythmic agents of Class I according to the classification scheme of Vaughan-Williams ("Classification of antiarrhythmic drugs" In: Cardiac Arrhythmias, edited by: E. Sandoe, E. Flensted-Jensen, K. Olesen; Sweden, Astra, Sodertalje, pp449-472, 1981) which cause a selective inhibition of the
20 maximum velocity of the upstroke of the action potential (V_{max}) are inadequate for preventing ventricular fibrillation. In addition, they have problems regarding safety, namely, they cause a depression of
25 myocardial contractility and have a tendency to induce arrhythmias due to an inhibition of impulse conduction. Beta-adrenoceptor blockers and calcium antagonists which belong to Class II and IV, respectively, have a defect in that their effects are either limited to a certain type of
30 arrhythmia or are contraindicated because of their cardiac depressant properties in certain patients with cardiovascular disease. Their safety, however, is higher than that of the antiarrhythmic agents of Class I.

Antiarrhythmic agents of Class III are drugs that cause a selective prolongation of the duration of the action potential without a significant depression of the V_{max} . Available drugs in this class are

- 2 -

limited in number. Examples such as sotalol and amiodarone have been shown to possess interesting Class III properties (Singh B.N., Vaughan Williams E.M. "A third class of anti-arrhythmic action: effects on atrial and ventricular intracellular potentials and other pharmacological actions on cardiac muscle, of MJ 1999 and AH 3747" *Br. J. Pharmacol.* 1970; 39:675-689. and Singh B.N., Vaughan Williams E. M, "The effect of amiodarone, a new anti-anginal drug, on cardiac muscle", *Br J. Pharmacol* 1970; 39:657-667.), but these are not selective Class III agents. Sotalol also possesses Class II effects which may cause cardiac depression and is contraindicated in certain susceptible patients. Amiodarone, also is not a selective Class III antiarrhythmic agent because it possesses multiple electrophysiological actions and is severely limited by side effects (Nademanee, K. "The Amiodarone Odyssey". *J. Am. Coll. Cardiol.* 1992; 20:1063-1065.) Drugs of this class are expected to be effective in preventing ventricular fibrillations. Selective class III agents, by definition, are not considered to cause myocardial depression or an induction of arrhythmias due to inhibition of conduction of the action potential as seen with Class I antiarrhythmic agents.

Class III agents increase myocardial refractoriness via a prolongation of cardiac action potential duration. Theoretically, prolongation of the cardiac action potential can be achieved by enhancing inward currents (i.e. Na^+ or Ca^{2+} currents; hereinafter I_{Na} and I_{Ca} respectively) or by reducing outward repolarizing potassium (K^+) currents. The delayed rectifier (I_{K}) K^+ current is the main outward current involved in the overall repolarization process during the action potential plateau, whereas the transient outward (I_{to}) and inward rectifier (I_{K1}) K^+ currents are responsible for the rapid initial and terminal phases of repolarization, respectively. Cellular electrophysiologic studies have demonstrated that I_{K} consists of two pharmacologically and kinetically distinct K^+ current subtypes, I_{Kr} (rapidly activating and deactivating) and I_{Ks} (slowly activating and deactivating). Class III antiarrhythmic agents currently in development, including d-sotalol, dofetilide (UK-68,798), almokalant (H234/09), E-

- 3 -

4031 and methanesulfonamide-N-[1'-6-cyano-1,2,3,4-tetrahydro-2(R)-naphthalenyl]-3,4-dihydro-4(R)-hydroxyspiro[2H-1-benzopyran-2,4'-piperidin]-6-yl],(+)-,monochloride predominantly, if not exclusively, block I_{K_r} . Although, amiodarone is a blocker of I_{K_s} (Balser J.R. Bennett, P.B., Hondeghem, L.M. and Roden, D.M. "Suppression of time-dependent outward current in guinea pig ventricular myocytes: Actions of quinidine and amiodarone. *Circ. Res.* 1991, 69:519-529), it also blocks Na^+ and Ca^{2+} currents, effects thyroid function, is as a nonspecific adrenergic blocker, and acts as an inhibitor of the enzyme phospholipase (Nademanee, K. "The Amiodarone Odessey". *J. Am. Coll. Cardiol.* 1992; 20:1063-1065). Therefore, its method of treating arrhythmia is uncertain.

Most Class III agents that are known to be in development predominantly block I_{K_r} . These agents have a potential liability in that they have an enhanced risk of proarrhythmia at slow heart rates. For example, torsades de points has been observed when these compounds are utilized (Roden, D.M. "Current Status of Class III Antiarrhythmic Drug Therapy", *Am J. Cardiol.* 1993; 72:44B-49B). This exaggerated effect at slow heart rates has been termed "reverse frequency-dependence", and is in contrast to frequency-independent or frequency-dependent actions (Hondeghem, L.M. "Development of Class III Antiarrhythmic Agents". *J. Cardiovasc. Cardiol.* 20 (Suppl. 2):S17-S22).

A number of antiarrhythmic agents have been reported in the literature, such as those disclosed in:

- (1) EP-A-O 397,121-A,
- (2) EP-A-O 300,908-A,
- (3) EP-A-O 307,121,
- (4) U.S. 4,629,739,
- (5) U.S. 4,544,654,
- (6) U.S. 4,788,196,
- (7) EP application No. 88302597.5,
- (8) EP application No. 88392598.3,
- (9) EP application No. 88302270.9,

- 4 -

- 5 (10) EP application No. 88302600.7,
(11) EP application No. 88302599.1,
(12) EP application No. 88300962.3,
(13) EP-A-O 235,752,
(14) DE No. 3633977-A1,
(15) U.S. 4,804,662,
(16) U.S. 4,797,401,
(17) U.S. 4,806,555,
(18) U.S. 4,806,536,
10 (19) U.S. 5,032,598,
(20) U.S. 5,032,604.

Despite considerable progress there is still a need for new methods to treat cardiac arrhythmia. In an effort to overcome the side effects
15 inherent in currently available antiarrhythmic agents, compounds that treat arrhythmia through blockade of I_{Ks} are presented.

OBJECTS OF THE INVENTION

It is, accordingly, an object of the present invention to
20 provide methods of treating cardiac arrhythmia with compounds not previously known to have activity for this condition. Another object is to provide new methods for treating cardiac arrhythmia. A further object is to provide pharmaceutical formulations and methods for their preparation for use in treating cardiac arrhythmia. These and other
25 objects of the present invention will be apparent from the following description.

SUMMARY OF THE INVENTION

A method of treating cardiac arrhythmia in mammals
30 comprising block of the slowly activating delayed rectifier potassium (K^+) current (I_{Ks}) is presented.


Among the compounds which exemplify this method of treatment are the 1,4-benzodiazepines or benzodiazepine derivatives that

- 5 -



block the I_{Ks} current and are therefore effective in the treatment of cardiac arrhythmia.

Additionally, it has been found that compounds, which at a concentration of 1 μ M or less (IC_{50}), selectively block 50% of the I_{Ks} current measured in isolated myocytes and exhibit a selectivity ratio equal to or greater than 10 times the block of the I_{Kr} , I_{K1} currents, result in treatment which prolongs the cardiac refractory period equally well or to a greater extent as heart rate increases, producing a frequency-independent or frequency-dependent treatment.

Benzodiazepine analogs of the general formula




wherein A is a 6-membered saturated or unsaturated carbocyclic ring or a 6-membered heterocyclic ring containing N, or N and O, Y is $-NH_2$, $NHSO_2R^1$,



wherein R^1 is

30

- 6 -



10 n is 0 or 1, X is O or -CH₂R, R² is R or -CONHSO₂R and R is straight or branched C₁₋₆ alkyl or C1-3alkylamine wherein the amino group is optionally mono- or di-substituted by C₁₋₃alkyl have been found to be useful in treating cardiac arrhythmia.

15 DETAILED DESCRIPTION

A method of treating cardiac arrhythmia in mammals comprising block of the slowly activating delayed rectifier potassium (K⁺)current (I_{Ks}) is presented.

20 Among the compounds which exemplify this method of treatment are the 1,4-benzodiazepines or benzodiazepine derivatives that block the I_{Ks} current and are therefore effective in the treatment of cardiac arrhythmia.

25 Additionally, it has been found that compounds, which at a concentration of 1 μM or less (IC₅₀), selectively block 50% of the I_{Ks} current measured in isolated myocytes and exhibit a selectivity ratio equal to or greater than 10 over block of the I_{Kr}, I_{K1} currents, result in treatment which prolongs the cardiac refractory period equally well or to a greater extent as heart rate increases, producing a frequency-independent or frequency-dependent treatment.

30 Several of the compounds used according to the present invention are known per se as antagonists of gastrin and cholecystikinin (CCK). The invention includes racemic mixtures of these compounds, the separated R and S enantiomers, and preferably, the R enantiomer. Examples of suitable compounds according to the present invention are

- 7 -

disclosed in US Patent 4,820,834, EP-A-0 434 360 and EP-A-0 434 369. Examples of some preferred compounds are listed below of this type:

- 5 1. 3(R)-(+)-1,3-dihydro-3-(2-indolecarbonylamino)-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one.
2. 3(S)-(-)-1,3-dihydro-3-(2-indolecarbonylamino)-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one.
- 10 3. N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea.
4. (R)-N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea.
- 15 5. N-[3(R,S)-5-cyclohexyl-2,3-dihydro-1-methyl-2-oxo-1H-1,4-benzodiazepin-3-yl]-N'-[3-(isopropylsulfonylaminocarbonyl)phenyl]urea, example 2 herein.
- 20 6. N,N-dimethyl-4-(3(R,S)-(((3-methylphenyl)amino)-carbonyl)amino-1,3-dihydro-1-(2-methylpropyl)-2-oxo-1,4-benzodiazepin-5-yl)phenylmethanamine.
- 25 7. (R)-3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one.
8. N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(m-methylphenyl)urea.
- 30 9. (+-)-N-(2,3-dihydro-1-methyl-2-oxo-5-(4-pyridinyl)-1H-1,4-benzodiazepin-3-yl)-1H-indole-2-carboxamide.

- 8 -

10. N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-4-methylbenzenesulfonamide.

11. (R)-1-(2-(dimethylaminoethyl)-5-(2-fluorophenyl)-1,3-dihydro-3-((1-methyl-1H-indole-3-yl)methyl)-2H-benzodiazepin-2-one.

12. N-(2,3-dihydro-1-methyl-2-oxo-5-(4-morpholino)-1H-1,4-benzodiazepin-3-yl)-N'-3-methylphenylurea.

Additional benzodiazepines or benzodiazepine derivatives have also been found to block the IK_s current. Representative compounds capable of blocking this current are presented in the Example section below

Certain compounds have been shown to exhibit selective block of the IK_s current. Examples of these compounds are:

a. (3RS)-N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)-urea.

b. (3R)-N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)-urea.

c. E-(+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)-urea.

d. (+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(3-chlorophenyl)propanamide.

e. (3RS)-3-cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl)propanamide.

f. (-)-3-Carbohexyl-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-4-oxido-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide.

- 9 -

- g. (+)-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-hexanamide.
- 5 h. (+)-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-pentanamide.
- i. N-[2,3-Dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-cyclohexylpropanamide.
- 10 j. N-[2,3-dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide.
- 15 k. E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(4-methoxyphenyl)-2-propenamide.
- l. (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide.
- 20 m. (+)-3-Cyclohexyl-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide.
- 25 n. (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-(2,4-dichlorophenylthio)-acetamide.
- 30 o. (+)-N-[(3R)-7-Amino-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide.
- p. (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-4-phenylbutanamide.

- 10 -

- q. (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-5-methyl-3-phenylisoxazole-4-carboxamide.
- 5 r. N-[2,3-Dihydro-1-methyl-2-oxo-5-(4-methoxyphenyl)-1H-1,4-benzodiazepin-3-yl]-3-[2,4-dichlorophenyl]-propanamide.
- 10 s. N-[2,3-Dihydro-1-methyl-2-oxo-5-ethyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide.
- t. E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)-2-propeneneamide

15

The utility of the exemplified compounds as anti-arrhythmic agents is shown by the following voltage-clamp and action potential studies in guinea pig isolated ventricular myocytes *in vitro* and by studies in conscious and anesthetized dogs *in vivo*. Although several
20 of these compounds exhibit CCK antagonist activity, their antiarrhythmic activity stems primarily from block of the I_{K_S} current.

By block of the the I_{K_S} current is meant a reduction in the amplitude of the time-dependent net outward current as measured in an isolated guinea pig myocyte and elicited during voltage clamp of the cell
25 from a holding potential of -50 mV to +50 mV for 1 second under the conditions defined in the methods in Section I below. Current amplitude is measured as the difference from the initial instantaneous current level following the capacitance discharge after changing the transmembrane voltage from -50 mV to +50 mV, to the current level at
30 the end of the 1 second voltage step. Normally, an increase in the current amplitude occurs as a function of time and voltage during depolarization of the cardiac cell membrane and this increasing current has been defined as the delayed rectifier K^+ current (I_K). Cellular electrophysiologic studies have demonstrated that I_K consists of two

- 11 -

pharmacologically and kinetically distinct K^+ current subtypes, I_{K_r} (rapidly activating and deactivating) and I_{K_s} (slowly activating and deactivating). I_{K_s} current occurs as a consequence of time- and voltage-dependent opening of ionic channels or pores in the cardiac cell membrane that allow the relatively selective flow of potassium ions (K^+) from the inside to the outside of the cell, that is, down the voltage and concentration gradient for K^+ ions that is inherent in cardiac cells. Thus, block of the I_{K_s} current ultimately reflects block of certain ion channels or pores in the cardiac cell membrane. However, the single pores or channels thought to underlie the I_{K_s} current have yet to be measured or identified at the single channel level.

By selective block of I_{K_s} is meant, compounds which block the I_{K_s} current measured in isolated myocytes by 50% at a concentration of $1\mu M$ or less (IC_{50}) and the concentration that blocks I_{K_s} by 50% is at least 10 fold lower than the concentration required to cause 50% block of I_{K_r} and/or I_{K_1} .

I. VOLTAGE CLAMP MEASUREMENT OF IONIC CURRENTS AND ACTION POTENTIALS *IN VITRO*

Cell preparation:

Guinea pig ventricular myocytes were isolated using a modification of the procedure described by Mitra *et al.*, *Am. J. Physiol.* 249:H1056-60 (1985). Excised hearts were perfused through the aorta (retrograde fashion) at a rate of 10 ml/min with oxygenated, warmed ($37^\circ C$) solutions described below. The heart was initially perfused for 7 minutes with a Tyrode's solution containing, in mM: 132 NaCl, 4 KCl, 1.2 $MgCl_2$, 10 HEPES, 5 glucose, pH = 7.2. The heart was then perfused with the same nominally Ca^{2+} -free solution containing 150 units/ml Type II collagenase (Worthington) and 0.5 units/ml Type XIV protease (Sigma) for 8 minutes. This perfusate was followed by a Tyrode's solution containing 0.2 mM $CaCl_2$ (without enzymes) for 5 minutes. The digested ventricles were cut into small pieces and gently

- 12 -

shaken until cells were visibly dispersed. The cells were stored at room temperature until use, within 8 hours after isolation.

Microelectrode fabrication:

5 The suction microelectrode technique described by Giles, W.R. and Shibata, E.F., 'Voltage clamp of bull-frog cardiac pacemaker cells: a quantitative analysis of potassium currents'. *J. Physiol.* 368:265-292 (1985) was used to voltage clamp the cells. Microelectrodes were made by using a two stage puller from square bore (1.0 mm o.d.)
10 borosilicate capillary tubing. Pipettes were filled with 0.5 M K gluconate, 25 mM KCl and 5 mM K₂ATP. The electrodes had resistances of 3 to 7 Mohm when filled with this solution. After establishing whole-cell recording mode, negative pressure was maintained on the pipette using a 1 ml gas-tight syringe attached via air
15 tight tubing to the suction port of the microelectrode holder. This minimized dialysis of the cell with the pipette solution.

Voltage-clamp technique:

20 To record K⁺ currents, the cells were bathed in a Ca²⁺-free Tyrode's solution containing 0.4 mM nisoldipine. Nisoldipine is a relatively specific blocker of L-type Ca²⁺ channels, having no effect on K⁺ currents at this concentration (Kass, R.S., "Nisoldipine: a new, more selective calcium current blocker in cardiac Purkinje fibers"., *J. Pharmacol. Exp. Ther.* 223:446-456, 1982). To record Ca²⁺ currents,
25 the cells were bathed in Tyrode's solution containing 1.8 mM Ca²⁺ without nisoldipine. The cell chamber was superfused with solutions at a rate of 1-2 ml/minute, while maintaining the temperature at 35±1°C. A List EPC-7 clamp amplifier was used to voltage clamp the isolated cells. Series resistance was compensated 40-70%, and current was low-pass
30 filtered at a cut-off frequency of 1 kHz. Data acquisition and analysis were performed using pClamp software (Axon Instruments, Burlingame, CA) and either an AST 286 or a Compaq 386 computer. A variety of voltage pulse protocols were used to measure Ca²⁺ current and three types of K⁺ currents: inward rectifier (IK1), rapidly

- 13 -

activating delayed rectifier (I_{K_r}) and the slowly activating delayed rectifier (I_{K_s}). Sanguinetti M.C. and Jurkiewicz, N.K., "Two components of cardiac delayed rectifier K^+ current: Differential sensitivity to block by Class III antiarrhythmic agents". *J. Gen Physiol* 5 1990; 96:194-214. Data was analyzed as % block from control membrane currents.

Current-clamp technique:

To record action potentials, single cells were bathed in 1.8 mM Ca^{2+} -Tyrode's solution. The cell chamber was superfused with 10 solutions at a rate of 1-2 ml/minute, while maintaining the temperature at $35 \pm 1^\circ C$. A List EPC-7 clamp amplifier was used to current-clamp the cells. Action potentials were elicited at a frequency of 0.2 Hz with 1 nA current pulses of 2 msec duration. Data were analyzed as percent 15 (%) change in action potential duration at 90% of repolarization (APD_{90}).

Solutions:

Nisoldipine was prepared as a 4 mM stock solution in 20 polyethylene glycol 200. Compounds tested in this series were prepared as 10 mM stock solutions in polyethylene glycol 400 or dimethyl sulfoxide. At the final concentrations used neither polyethylene glycol 200, 400 nor dimethyl sulfoxide had any effect on membrane currents.

Results:

As an example of this series of compounds, compound number 4 in the foregoing list of preferred compounds exhibited a concentration dependent block of I_{K_s} with an IC_{50} of 215 nM. Complete block of I_{K_s} was observed at 5 mM. Little or no effect was 30 seen on I_{K_1} (1 mM), I_{K_r} (1 mM) or I_{Ca} (0.1 mM) at the indicated concentrations. IC_{50} for I_{K_r} was estimated to be approximately 5 mM. Action potentials elicited from 4 cells exposed to 5 mM of compound number 4 exhibited an increase in APD_{90} of $202 \pm 85\%$.

- 14 -

II. ELECTROCARDIOGRAPHIC QT INTERVAL IN CONSCIOUS DOGS

Method:

5 Trained, female mongrel dogs, in the weight range of 8 to 16 kg and instrumented with chronic arterial access ports, were deprived of food 18 to 24 hours prior to the study. At test time, the dog was placed in an Alice King Chatham nylon mesh sling for light restraint. Four Grass Instrument Co. Type E2 Platinum Subdermal
10 Electrodes were affixed to the limbs of the dog and connected to a Hewlett-Packard 4700A Cardiograph multi-lead electrocardiogram (ECG) recorder. After approximately a 30-minute stabilization period, one or two pretreatment Lead II ECG tracings were taken over approximately a 30-second interval. Dogs were then dosed via gavage
15 with labrafil (vehicle) or test compound. The volume administered was 2.0 ml/kg followed by a 15 ml tap-water flush. Lead II ECG recordings were obtained at 15, 30, and 60 minutes and every 30 minutes thereafter for five hours post-drug administration. Dogs were also monitored for blood pressure changes via a Modular Instruments
20 data acquisition system and transducer connected to the arterial access port. Final ECG traces were taken at approximately 24 hours post-treatment. Average QT interval lengths were determined for each time point and QT_c values were calculated to correct for variations in heart rate ($QT_c = QT/\sqrt{R-R}$).

25 As an example of this series of compounds, oral administration of compound number 4 elicited a dose-related prolongation of ECG QT interval. There were no other ECG or cardiovascular effects of compound number 4 in conscious dogs.

30 III. FREQUENCY-DEPENDENCE OF CLASS III ACTIVITY

The propensity for Class III antiarrhythmic agents that predominantly block IK_r to cause arrhythmia at slow heart rates is likely due to the "reverse frequency-dependent" action of these agents,

- 15 -

that is, these agents have greater activity at slow vs fast heart rates (Hondegheem, L.M. and Snyders, D.J., "Class III antiarrhythmic agents have a lot of potential, but a long way to go: reduced effectiveness and dangers of reverse use-dependence." *Circulation* 81:686-690, 1990; Hondegheem, L.M., "Development of Class III antiarrhythmic agents". *J. Cardiovasc. Pharmacol.* (Suppl 2): S17-S22, 1992; Funck-Brentano, C., "Rate-dependence of Class III actions in the heart". *Fundam. Clin. Pharmacol.* 7:51-59, 1993). Cellular electrophysiologic studies indicate that the reverse frequency-dependent profile Class III agents that block I_{K_r} is the result of a diminished contribution of the I_{K_r} current, relative to other outward currents (eg., I_{K_s}), to repolarization at fast heart rates (Jurkiewicz N.K. and Sanguinetti, M.S. "Rate-dependent prolongation of cardiac action potentials by a methanesulfonanilide Class III antiarrhythmic agent: specific block of rapidly activating delayed rectifier K^+ current by dofetilide" *Circ. Res.* 72:75-83, 1993). Because I_{K_s} may contribute more to repolarization at faster heart rates, the selective blockade of I_{K_s} represents a potential target for safer and more effective Class III antiarrhythmic agents. First, a selective blocker of I_{K_s} should not display exaggerated activity at slow heart rates, and would therefore possess a more favorable safety profile than reverse frequency-dependent I_{K_r} blocker Class III antiarrhythmic agents. Second, a selective blocker of I_{K_s} should possess equal or greater activity at faster heart rates which might result in greater efficacy against arrhythmias. The *in vitro* cellular electrophysiologic studies and *in vivo* cardiac electrophysiologic studies in anesthetized dogs described below were conducted in order to test the hypothesis that the selective blockade of I_{K_s} would result in either frequency-independent (i.e., equivalent activity at slow and fast rates) or forward frequency-dependent (greater activity at faster vs slower rates) Class III activity.

- 16 -

III. A. FREQUENCY-DEPENDENT EFFECTS ON ACTION POTENTIAL DURATION *IN VITRO* IN ISOLATED VENTRICULAR MYOCYTES

Methods:

5

Cell preparation: Same as described in section I.

Action Potential Measurements:

10 Transmembrane action potential measurements were made
using a technique similar to that described previously by Salata, J.J. and
Wasserstrom, J.A. "Effects of quinidine on action potentials and ionic
currents in isolated canine ventricular myocytes", *Circ. Res.* 62:324-337
(1988). The isolated ventricular myocytes were transferred to a glass
15 bottom experimental perfusion chamber (12x25 mm) filled with a
HEPES-Tyrode's solution containing in mM: 132 NaCl, 4 KCl, 1.8
CaCl₂, 1.2 MgCl₂, 10 HEPES, 5 glucose, pH = 7.2 to a height of 2 mm.

After the myocytes settled to the bottom of the chamber, they were
continuously superfused with preheated HEPES-Tyrode's solution at a
rate of 2.0 ml/min. The temperature was monitored and maintained at
20 36±0.5°C. The experimental chamber was placed on the stage of an
inverted microscope (Nikon Diaphot-TMD) to view the cells.

Transmembrane potentials were recorded by means of conventional
glass microelectrodes that were filled with 3 M KCl and had tip
resistances of 40-60 MΩ. The microelectrode was connected through a
25 Ag-AgCl electrode to the headstage preamplifier (model HS-2,
gain=1.0) of an operational amplifier with a high input impedance and
variable input capacitance neutralization (Axoclamp -2A, Axon
Instruments, Foster City, CA). The voltage drop across the
microelectrode produced by current injection was compensated
30 electronically with an active bridge circuit immediately before
impalement of a cell. The experimental bath was grounded through a 3
M KCl agar bridge and a Ag-AgCl junction. The microelectrodes were
positioned carefully at the cell surface using a hydraulic
micromanipulator (Narishige Model MO-303, Greenvale, NY).

- 17 -

Penetration of the cell membrane was achieved by briefly (1-3 msec) increasing the input capacitance neutralization to cause oscillation of the amplifier. After establishing a stable microelectrode seal, the myocyte was lifted off the bottom of the chamber to allow complete superfusion of the cell. Action potentials were generated by passing current pulses of 1 msec in duration, and with an intensity of 1.2 times threshold at varying stimulus rates through the recording electrode. Cells were allowed to stabilize for 20 minutes before collection of data began. Voltage signals were digitized using an analog to digital converter (TL-1, DMA interface, Axon Instruments, Foster City, CA) at a sampling frequency of 2 kHz. Data acquisition and analysis were performed with pClamp software (Axon Instruments, Foster City, CA) and an AST 486 computer (Irvine, CA). Action potentials were stimulated at rates of either 60, 120, and 180 beats/min and were allowed to stabilize for 1 minute at each rate. Trains of stimuli were applied at each of these rates for a duration of 30 seconds and the last action potential of the train was recorded from each of 5 trains and these 5 action potentials were averaged. Action potential duration at 90% of repolarization (APD₉₀) was measured from the averaged action potential traces at each stimulus rate in control and at 10 minutes after superfusion of solution containing the compound of Example 26 at a given concentration. The compound was superfused at sequentially increasing concentrations. The compound of Example 26 was dissolved at a stock concentration of 1 mM in dimethyl sulfoxide (DMSO) and diluted directly into HEPES Tyrode's solution using serial dilutions as necessary to achieve final test concentrations.

Results:

In these studies, one of the preferred compounds, the compound of Example 26, produced both concentration and rate-dependent increases in the action potential duration of isolated guinea pig myocytes as shown in Table 1. In the control, APD₉₀ values were 229±22, 206±15, and 189±10 msec at stimulus rates of 60, 120, and 180 beats/min, respectively. The compound of Example 26, at 10 nM,

- 18 -

increased APD90 more at fast vs slow rates. For example, APD90 was increased an average of 12.4% at 180 beats/min vs 9.4% at 60 beats/min. Thus, at lower concentrations the compound increased APD90 in a forward frequency-dependent manner. At higher concentrations, APD90 was increased equally at fast and slow rates. The percent increases were not statistically different at the different stimulus rates, therefore, at higher concentrations APD90 was increased in a frequency-independent manner. The increases in APD90 were nearly completely reversible after 20 min of washout of the compound.

TABLE 1

E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)-2-propeneneamide :

Concentration- and Frequency-Dependent Increases in Action Potential Duration at 90% of Repolarization (APD90) in Isolated Ventricular Myocytes

- 19 -

STIMULUS

	Concentration [nM]	Rate beats/min	APD90 % Change	SEM
5	10	60	9.4	±2.4
		120	10.1	±1.2
		180	12.4	±3.4
10	30	60	16.0	±2.3
		120	14.8	±1.3
		180	14.4	±1.8
15	100	60	24.7	±4.4
		120	24.3	±4.4
		180	22.9	±4.0
20	Washout	60	4.2	±6.3
		120	5.3	±5.1
		180	5.9	±3.9

Data are Mean ± SEM (n≥5 cells)

25

III.B FREQUENCY- DEPENDENT EFFECTS ON CARDIAC REFRACTORY PERIOD *IN VIVO* IN ANESTHETIZED DOGS

Methods:

30

Purpose-bred male or female mongrel dogs (9.8-10.8 kg) were anesthetized with alpha chloralose (80 - 100 mg/kg i.v.), and the animals were ventilated using a volume-cycled respirator. The right femoral artery and vein were isolated and cannulated for the measurement of systemic arterial pressure and for test agent

- 20 -

administration, respectively. A right thoracotomy was performed in the 4th intercostal space, the pericardium incised, and the heart suspended in a pericardial cradle. One stainless steel bipolar plunge electrode was sutured to the anterior surface of the right ventricle for the determination of ventricular excitation threshold and refractory periods, while a second stainless steel bipolar plunge electrode was sutured to the right ventricular apex for ventricular pacing. A 21 gauge stainless steel hypodermic needle attached to a 1 ml syringe containing 40% formaldehyde was introduced into the right atrium via a stab wound and purse-string suture. The atrioventricular (AV) node was identified and then chemically ablated with formaldehyde using the method of Steiner and Kovalik (Steiner C. and Kovalik, "A simple technique for production of chronic complete heart block in dogs" T.W.; *J. Appl. Physiol.* 25:631-632, 1968). After the ablation of the AV node, heart rate was controlled and maintained at either 60 or 150 beats/min, as dictated by the experimental protocol, by ventricular apical pacing. Limb electrodes were attached for the continuous recording of Lead II electrocardiogram.

Ventricular excitation threshold (ET) and relative refractory periods (RRPs) were determined alternately at heart rates of 60 and 150 beats/min using an extrastimulus technique described previously (Wallace A.A. *et. al.*, "Cardiac electrophysiologic and inotropic actions of new and potent methanesulfonanilide Class III antiarrhythmic agents in the anesthetized dog". *J. Cardiovasc. Pharmacol.* 18:687-695, 1991). Ventricular excitation threshold was defined as the minimum current required to evoke a propagated response for a ventricular extrastimulus introduced at a coupling interval of 250-300 msec after a basic ventricular apical beat. The ventricular relative refractory period was defined as the longest coupling interval failing to elicit a propagated response for a ventricular extrastimulus introduced with a current intensity of 2x the ventricular excitation threshold. Multiple alternating baseline 60 and 150 beat/min RRP's were measured in each preparation, with two-to-three stable

- 21 -

consecutive baseline RRP determinations required prior to administration of the test agent.

In this study, one of the preferred compounds, E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)-2-propeneneamide, (Compound of Example 26) was used as an example of a selective blocker of I_{Ks} . The compound of Example 26 was administered intravenously at a dosage of 1.0 mg/kg over a period of 5 minutes to a group of 5 anesthetized dogs. The compound of Example 26 was infused in a vehicle of 2% cremophor/2% ethanol/96% saline, at a test agent concentration of 0.5 mg/ml. Ventricular RRP's at 60 and 150 beats/min were redetermined immediately, 15 and 30 minutes following the intravenous administration of the compound of Example 26. The frequency-dependence of the activity of the compound of Example 26 was assessed by comparing absolute increases in ventricular RRP's from the final corresponding baseline value baseline determined at slower (60 beat/min) vs faster (150 beat/min) heart rates.

Results:

The table below summarizes the absolute increases in the ventricular RRP from the respective baseline values observed at slower (60 beat/min) vs faster (150 beat/min) heart rates after the intravenous administration of 1.0 mg/kg of the compound of Example 26. Equivalent increases in the ventricular RRP's were observed at the 60 and 150 beat/min heart rates following the administration of the compound of Example 26, indicating that it displays "frequency-independent" activity to increase the RRP.

TABLE 2

Increase in Ventricular Relative Refractory Period (msec) Above Baseline Following the Intravenous Administration of 1.0 mg/kg of the compound of Example 26 at Indicated Heart Rate and Time Points after Infusion

- 22 -

	Heart Rate (Beat/Min)	Immediate	15 Min	30 Min
5	60	29.6±4.4	19.2±3.4	14.0±2.3
	150	23.2±7.3	15.6±4.1	11.4±4.4

Data are Mean±S.E.M.; n=5

10

The compounds of the present invention have the pharmacological properties required for the antiarrhythmic agents of Class III, namely the prolongation of the myocardial action potential *in vitro* without a significant depression of the V_{max} or prolongation of cardiac refractory period or QTC-interval in dogs. Moreover, the effects of many of the novel compounds, in contrast to such I_{Kr} blockers, increase action potential duration equally or to a greater extent at fast heart rates. These compounds are much more potent than the reference drug, sotalol.

15

20

The compounds of the present invention are effective in treating and preventing all types of arrhythmias including ventricular and atrial (supraventricular) arrhythmias. The compounds of the present invention are especially useful to control reentrant arrhythmias and prevent sudden death due to ventricular fibrillation.

25

In the novel method of treating arrhythmia of this invention, a compound or pharmaceutically acceptable salt thereof, is administered in an amount ranging from about 0.1 to about 50 mg per kg of body weight per day, preferably from about 1.0 to about 30 mg per kg of body weight per day in a single dose or in 2 to 4 divided

30

doses. The compounds of this invention can be administered as the sole active ingredient or in combination with other antiarrhythmic agents or other cardiovascular agents.

- 23 -

The compounds, or pharmaceutically acceptable salts thereof, of the present invention, in the described dosages, are administered orally, intraperitoneally, subcutaneously, intramuscularly, transdermally, sublingually or intravenously. They are preferably administered orally, for example in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gum, or the like prepared by art recognized procedures. The amount of active compound in such therapeutically useful compositions or preparations is such that a suitable dosage will be obtained.

The following examples illustrate the present invention without, however, limiting the same thereto.

EXAMPLE 1

Following the protocol described previously but substituting the compounds listed below for compound number 4, the following results are obtained:

	IC ₅₀ or % Block (μM)		IC ₅₀ (mM)	
	Guinea Pig Myocytes			
No. of Compound	IKs	IKr	CCKA	CCKB
1	0.220	20% (1)	8.0	3,800
2	25% (1)	21% (1)	0.07	220
5	57	77	>3mM	>3mM
			2%	0%
8	10	38	-	-
9	20	93	-	-

- 24 -

Cellular Electrophysiology (% Block)

		Conc.	I _{K1}	I _{Kr}	I _{Ks}
5	5	1 μ M	n = 2 1.6	n = 2 16.1	n = 2 10
		10 μ M	n = 5 5.2+1.8	n = 5 24.7+13	n = 5 30.5+6.6
	7	1 μ M	n = 4 6.5+2.	n = 4 14.8+6.7	n = 6 29.9+6
10		10 μ M	n = 4 8.6+2.2	n = 4 36.4+2.9	n = 4 43.4+10.8

15 Cellular Electrophysiology (% Block)

		Conc.	I _{K1}	I _{Kr}	I _{Ks}
20	9	1 μ M	n = 2 2	n = 2 44.3	n = 2 20
		10 μ M	n = 2 0.6	n = 2 82.4	n = 2 66
	12	1 μ M	n = 4 8+2.4	n = 4 13+3.1	n = 4 14+6.5
25		10 μ M	n = 4 10+4.2	n = 4 30+5.4	n = 4 16+11

EXAMPLE 2

30 N-[3(R,S)-5-Cyclohexyl-2,3-dihydro-1-methyl-2-oxo-1H-4-benzo-diazepin-3-yl]N'-[3-(isopropylsulfonyl amino carbonyl)phenyl]urea

Step 1: Isopropylsulfonamide

Ammonia gas was bubbled through a stirred solution of isopropylsulfonyl chloride (3.9 ml, 35 mmol) in anhydrous

- 25 -

tetrahydrofuran (100 ml), cooled to 0°C, for 30 minutes. After allowing to warm to ambient temperature, the mixture was filtered and the filtrate evaporated *in vacuo*, to leave a white solid. This was partitioned between ethyl acetate (50 ml) and water (50 ml). The organic phase was separated and the aqueous phase extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried (Na₂SO₄) and evaporated *in vacuo* to afford the title compound (3.5 g, 81%) as a colorless solid, mp 51-53°C. ¹H NMR (360 MHz, CDCl₃) w 1.92 (6H, d, J = 6.7 Hz), 3.22 (1H, septet, J = 6.7 Hz), 4.61 (2H, br s).

Step 2: 1-(Isopropylsulfonylaminocarbonyl)-3-nitro-benzene

To a mixture of isopropylsulfonamide (1.7 g, 13.8 mmol), 3-nitrobenzoic acid (2.31 g, 13.8 mmol) and 4-dimethylaminopyridine (1.69 g, 13.8 mmol) in anhydrous dichloromethane (100 ml) under an atmosphere of nitrogen was added 1-[3-(dimethylamino)-propyl]-3-ethyl carbodiimide hydrochloride (2.65 g, 13.8 mmol). The mixture was stirred at ambient temperature for 20 hours, then extracted with 1M NaOH and the separated aqueous phase was acidified using 5M HCl. The solid which precipitated was collected by filtration, washed with water and dried under vacuum to afford the title compound (2.74 g, 75%) as a colorless solid, mp 175-177°C. ¹H NMR (360 MHz, D₆-DMSO) w 1.34 (6H, d, J = 6.9 Hz), 3.83 (1H, septet, J = 6.9 Hz), 7.93 (1H, dd, J = 8.0 and 8.0 Hz), 8.35 (1H, d, J = 8.0 Hz), 8.48 (1H, d, J = 8.0 Hz), 8.78 (1H, s), 12.40 (1H, br s).

Step 3: 1-(Isopropylsulfonylaminocarbonyl)-3-amino-benzene

To a suspension of 1-(isopropylsulfonylaminocarbonyl)-3-nitrobenzene (2.5 g, 9.2 mmol) in ethanol (50 ml) was added 10% palladium on carbon (0.25 g, 10% (w/w)) in water (2 ml). The mixture was hydrogenated at 40 psi for 10 minutes after which the catalyst was filtered off and washed with ethanol. The solvents were evaporated *in vacuo* to give the title compound as a yellow solid. This was recrystallised from hot ethanol to give a pale yellow crystalline solid (1.7 g 75%), mp 190-193°C. ¹H NMR (360 MHz, D₆-DMSO) w

- 26 -

1.30 (6H, d, J = 6.8Hz), 3.79 (1H, septet, J = 6.9 Hz), 5.36 (2H, br s), 6.79 (1H, dd, J = 7.9 and 1.2Hz), 7.05 (2H, m), 7.13 (1H, dd, J = 7.8 and 7.8Hz).

5 Step 4: N-3-(R,S)-5-Cyclohexyl-2,3-dihydro-1-methyl-oxo-1H-1,4-benzodiazepin-3-yl] N'-[3-(isopropylsulfonylamino-carbonyl)phenyl]urea


 A solution of 5-cyclohexyl-1,3-dihydro-1-methyl-3(R,S)-[4-nitrophenyloxycarbonyl]-amino]-2H-1,4-benzodiazepin-2-one (0.3 g, 0.69 mmol) in anhydrous dimethyl formamide (6 ml) and 1-(isopropyl-carbonylaminosulfonyl)-3-aminobenzene (0.175 mg, 0.72 mmol) was added dropwise. After recrystallization from hot ethanol the compound (0.24 g, 65%) was afforded as a colorless solid, mp 165°C (dec.). ¹H NMR (360MHz, D₆-DMSO) w 0.87-0.98 (1H, m), 1.10-1.64 (7H, m), 1.30 (6H, d, J = 6.9Hz), 1.79 (1H, m), 1.88-1.96 (1H, m), 2.95 (1H, m), 3.33 (3H, s), 3.79 (1H, septet, J = 6.9Hz), 5.07 (1H, d, J = 8.2Hz), 7.37 (3H, m), 7.46 (1H, d, J = 7.8Hz), 7.55 (2H, m), 7.64 (1H, dd, J = 7.1 and 7.1Hz), 7.75 (1H, d, J = 7.9Hz), 7.92 (1H, s), 9.22 (1H, s), 11.93 (1H, br s).

20

EXAMPLE 3








X Other compounds which exemplify the treatment of arrhythmia through blockade of the IKs channel include, but are not limited to the following

25



I

- 27 -

	R ¹	R ²	R ³	R ⁴	R ⁵
5		$-(CH_2)_2-\underset{\begin{array}{c} CH_3 \\ \end{array}}{C}-(CH_2)_2-$		-CH ₃	H
		$-CH_2-CH$  $CHCH_2-$		-CH ₃	H
10		$-CH_2-CH$  $CHCH_2-$		-CH ₃	H
15		-CH ₃	cyclohexyl	-CH ₃	H
20		$-(CH_2)_2-\underset{\begin{array}{c} \\ CH_3 \end{array}}{CH}-(CH_2)_2-$		-CH ₃	H

N-[3(RS)-5-(N-Cyclohexyl-N-methylamino)-2,3-dihydro-1-methyl-2-oxo-1H-1,4-benzodiazepin-3-yl]-N'-[3-methylphenyl]urea

Step A: Preparation of Methyl-2-(N-bromoacetyl-N-methylamino) benzoate

A solution of bromoacetyl bromide (209 g, 1.03 mol) in dichloromethane (200 ml) was added dropwise to a cooled (ice bath) solution of methyl N-methylantranilate (158 g, 0.96 mol) in dichloromethane (1.41). A solution of sodium hydroxide (59 g, 1.47 mol) in water (400 ml) was added dropwise to this ice cold solution then after addition the reaction mixture was stirred at room temperature for 20 hours. The organic phase was separated and washed with 1M hydrochloric acid (500 ml), brine (300 ml), saturated sodium hydrogen carbonate solution (400 ml), dried (sodium sulphate) then evaporated to

- 28 -

afford the required product as a solid (255 g, 92%). ¹H NMR (360MHz, CDCl₃) δ 3.23 (3H, s), 3.54 (1H, d, J = 11Hz), 3.60 (1H, d, J = 11Hz), 3.90 (3H, s), 7.40 (1H, d, J = 8Hz), 7.51 (1H, dd, J₁=J₂=8Hz), 7.65 (1H, dd, J₁=J₂=8Hz), 8.04 (1H, d, J = 8Hz).

5

Step B: Preparation of 2,5-Dioxo-1-methyl-1,4-benzodiazepine

Ammonia gas was bubbled through an ice-cooled solution of methyl 2-(N-bromoacetyl-N-methylamino)benzoate (255 g, 0.89 mol) in methanol (1.6l) until saturated. The cooling bath was removed and the reaction mixture left standing at room temperature for 18 hours. The precipitate was collected to afford the required product (79 g). The filtrate was evaporated and the residue partitioned between dichloromethane (300 ml) and 10% citric acid solution (200 ml). The organic layer was separated, washed with brine (200 ml), dried (sodium sulphate) then evaporated to give a solid which was recrystallized from dichloro-methane/petroleum ether (60-80) to afford further product (32.5 g). Total yield = 111.5 g (73%). Mp 190-193°C. ¹H NMR (360 MHz, CDCl₃) δ 3.42 (3H, s), 3.80 (2H, broad s), 6.80 (1H, s), 7.24 (1H, d, J = 8 Hz), 7.32 (1H, dd, J₁ = J₂ = 8 Hz), 7.57 (1H, dd, J₁ = J₂ = 8 Hz), 7.90 (1H, d, J = 8 Hz). Found C, 63.20; H, 5.25; N, 14.77. C₁₀H₁₀N₂O₂ requires C, 63.15; H, 5.30; N, 14.73%.

15
20

Step C:

The desired product was prepared as for Example 1 parts a, b, and c, using 2,5-dioxo-1-methyl-1,4-benzodiazepine in place of 2,5-dioxo-1-propyl-1,4-benzodiazepine to give a white solid which was recrystallized from methanol, water to give the desired product (70 mg). Mp 145-147°C. ¹H NMR (360 MHz, D₆-DMSO) δ 0.99 (10H, m), 2.21 (3H, s), 2.65 (3H, s), 3.31 (4H, m), 4.92 (1H, d, J = 8.4 Hz), 6.70 (1H, d, J = 8.0 Hz), 6.93 (1H, d, J = 7.5 Hz), 7.09 (2H, m), 7.17 (1H, s), 7.36 (1H, m), 7.49-7.69 (3H, m), 8.80 (1H, s); MS (CI) m/e 434 [MH]⁺. Anal. Found C, 67.22; H, 7.23; N, 15.29. C₂₅H₃₁N₅O₂·0.75 H₂O requires C, 67.17; H, 7.33; N, 15.67%.

25
30

- 29 -

EXAMPLE 4(-)-N-[5-(N-Cyclohexyl-N-methylamino)-2,3-dihydro-2-oxo-1-methyl-1H-1,4-benzodiazepin-3-yl]-N'-[3-methylphenyl]urea Hydrochloride

5 The desired product was separated from the racemate of Example 4 by chiral HPLC as for Example 2, Step A. Peak A (eluted first) was treated as for Example 2 part B and the free base obtained was dissolved in dichloromethane. Ethereal hydrogen chloride was added and after 5 minutes the solvent was removed *in vacuo*. The
10 resulting oil was crystallized from dichloromethane/ether to give the desired hydrochloride as a white solid (100 mg). Mp 193-195°C. ¹H (360MHz, D₆-DMSO, trifluoroacetic acid) δ 0.95-1.95 (10H, m), 2.23 (3H, s), 3.12 (3H, s), 3.43 (4H, m), 5.39 (1H, m), 6.76 (1H, d, J = 7.2Hz), 7.11-7.90 (7H, m). MS (CI) m/e [MH]⁺. Anal. Found. C, 60.45; H, 7.02; N, 14.05. C₂₅H₃₁N₅O₂·HCl. 1.5 H₂O requires C, 60.41; H, 7.02; N, 14.35%, [α]_D²² -195° (c=0.1, MeOH). Purity A:B
15 =>99%.

EXAMPLE 5


20 (+)-N-[5-(N-Cyclohexyl-N-methylamino)-2,3-dihydro-2-oxo-1-methyl-1H-1,4-benzodiazepin-3-yl]-N'-[3-methylphenyl]urea Hydrochloride

 The desired product was separated from the racemate from Example 4 by chiral HPLC as for Example 2 part A. Peak B (eluted
25 second) was treated as for Example 5 peak A to yield the desired hydrochloride (90 mg). Mp 194-196°C. ¹H NMR (360MHz, D₆-DMSO + trifluoroacetic acid) δ 0.95-1.95 (10H, m), 2.24 (3H, s), 3.12 (3H, s), 3.43 (4H, m), 5.39 (1H, m), 6.76 (1H, d, J = 7.5Hz), 7.06-7.86 (7H, m), 9.20 (1H, s); MS (CI) m/e 434 [MH]⁺. Anal. Found. C, 58.28; H, 6.82; N, 13.47. C₂₅H₃₁N₅O₂·HCl.2.35H₂O requires C, 58.61; H, 7.22; N, 13.67%. [α]_D²² +154° (C=0.1, MeOH). Purity B:A =>95%.

30

- 30 -

Further examples of other compounds which are useful in the treatment of arrhythmia by blockade of the IKs current include but are not limited to compounds such as:




where X is -CH₂-, R¹ is 1-naphtyl, R² and R³ taken together are -(CH₂)₆- and R⁴ is n-propyl.

EXAMPLE 6

(±)-N-[2,3-dihydro-1-methyl-2-oxo-5-(N-cyclohexyl-N-methylamino)-1H-1,4-benzodiazepin-3-yl]-3-phenylpropanamide

Step A:




A solution of PCl₅ (6.6 g, 32 mmol) in 250 mL dichloromethane was added to a stirred solution of 1-methyl-1,4-benzodiazepine-2,5-dione (5.0 g, 26 mmol) in 150 mL of dichloromethane. The solution was stirred at room temperature for 3 hours before

- 31 -

evaporation of volatiles. The resulting foam was dissolved in 200 mL dichloromethane, the solution cooled to 0°C and a solution of N-methylcyclohexylamine (11.8 mL, 91 mmol) in 50 mL of dichloromethane added over 5 minutes. The reaction mixture was allowed to warm to room temperature, and partitioned. The organic phase was washed with brine, dried (MgSO₄) and solvent evaporated to give the product as a foam. Yield 6.9 g.

NMR (300 MHz, CDCl₃) δ : 7.60 (m, 1H), 7.47 - 7.52 (m, 2H), 7.33 (m, 1H), 4.0 (1/2AB, J = 12.2 Hz, 1H), 3.47 (1/2AB, J = 12.2 Hz, 1H), 3.35 (s, 3H), 3.3 (m, 1H), 2.78 (s, 3H), 1.0-2.0 (m, 10H).

Step B:



A solution of 2,3-dihydro-1-methyl-2-oxo-5-(N-cyclohexyl-N-methylamino)-1H-1,4-benzodiazepine (770 mg, 2.7 mmol) in 10 mL toluene was added to a stirred and cooled (30°C) suspension of potassium t-butoxide (750 mg, 6.7 mmol) in 25 mL of toluene. After stirring at -30°C for 30 minutes, isoamyl nitrite (540 μ L, 4.0 mmol) was added and the reaction mixture stirred for 3 hours at -20°C. The mixture was then poured into 10% citric acid solution/ethyl acetate, stirred for 10 minutes, the pH adjusted to 7 with saturated potassium carbonate solution and the phases separated. The organic phase was washed with brine, the organic phase dried (MgSO₄) and the solvent evaporated to give a foam. This was dissolved in 15 mL THF and ethyl isocyanate (395 μ L, 5 mmol) added followed by

- 32 -

triethylamine (700 μ L, 5 mmol). The reaction mixture was heated to 60°C for 2 hours, cooled to room temperature, the volatiles evaporated and the residue purified by flash column chromatography (silica, 75% ethyl acetate/hexane) to afford 720 mg of product as a foam.

5

NMR (300 MHz, CD_3OD) δ : 7-7.6 (m, 5H), 3.5 (m, 1H), 3.42 (s, 3H), 2.7-3.3 (m, 5H), 1.1-2 (m, 10H), 1.05 (t, $J = 7$ Hz, 3H).


Step C:

10

15

20

25



30

A solution of the oxime carbamate (355 mg, 1.85 mmol) in 30 mL methanol was hydrogenated at 50 psi over 300 mg of 10% palladium/charcoal for 3 hours. The mixture was filtered through celite and the filtrate evaporated to give the crude amine. This was dissolved in 5 mL DMF and phenylpropionic acid (300 mg, 2 mmol), 1-hydroxybenzotriazole hydrate (305 mg, 2 mmol), triethylamine (250 μ L, 1.8 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (380 mg, 2 mmol) added. The reaction mixture was stirred at room

- 33 -

temperature for 18 hours, poured into water and extracted with ethyl acetate. The organic phase was washed with sodium bicarbonate solution and brine, dried (MgSO₄) and solvent evaporated to give a solid, recrystallized from ethyl acetate/hexane to afford (\pm)-N-[2,3-Dihydro-1-methyl-2-oxo-5-(N-cyclohexyl-N-methylamino)-1H-1,4-benzo-diazepin-3-yl]-3-phenylpropanamide.

m.p. 167-168°C.

Anal. Calcd. for C₂₆H₃₂N₄O₂•0.75 cyclohexane:

10 C, 73.9; H, 8.34; N, 11.3.

Found: C, 73.75; H, 8.36; N, 10.79%.

NMR (300 MHz, DMSO) δ : 8.8 (d, J = 7.5 Hz, 1H), 7.3-7.65 (m, 9H), 4.95 (d, J = 7Hz, 1H), 3.57 (1/2AB, J = 13 Hz, 1H), 3.48 (1/2AB, J = 13
15 Hz, 1H), 3.1-3.4 (m, 1H), 3.27 (s, 3H), 2.65 (s, 3H), 0.9-1.9 (m, 10H).

By substituting the appropriate acid for phenylpropionic acid, and employing the procedures substantially as described in Example 1 the following compounds were prepared.

20

EXAMPLE 7

(\pm)-N-[2,3-Dihydro-1-methyl-2-oxo-5-(N-cyclohexyl-N-methylamino)-1H-1,4-benzo-diazepin-3-yl]-2-cyclohexylacetamide

25

m.p. 158-159°C.

Anal. Calcd. for C₂₅H₃₆N₄O₂ 0.85 cyclohexane:

C, 72.86; H, 9.39; N, 11.29.

Found: C, 72.46; H, 9.4; N, 10.9%.

30 NMR (300 MHz, DMSO) δ : 8.44 (d, J = 7.8 Hz, 1H), 7.5-7.65 (m, 2H), 7.46 (dd, J = 7.8, 1.2 Hz, 1H), 7.32 (m, 1H), 4.95 (d, J = 8 Hz, 1H), 3.15-3.45 (m, 1H), 3.26 (s, 3H), 2.64 (s, 3H), 1.95-2.1 (m, 2H), 0.8-1.9 (m, 21H).

- 34 -

EXAMPLE 8

X
N-[2,3-Dihydro-1-methyl-2-oxo-5-(N-cyclohexyl-N-methylamino)-
1H-1,4-benzo-diazepin-3-yl]-4-phenylbutanamide

5

m.p. 140-141°C.

Anal. Calcd. for C₂₇H₃₄N₄O₂ 0.7 Water:

C, 70.62; H, 7.77; N, 12.2.

Found: C, 70.58; H, 7.54; N, 12.11%.

10

NMR (300 MHz, DMSO) δ: 8.52 (d, J = 8.1 Hz, 1H), 7.1-7.6 (m, 9H),
4.97 (d, J = 9.1 Hz, 1H), 3.1-3.45 (m, 1H), 3.26 (s, 3H), 2.64 (s, 3H),
2.52 (m, 2H), 2.10-2.30 (m, 2H), 0.9-1.9 (m, 12H).

15

EXAMPLE 9

(±)-N-[2,3-Dihydro-1-methyl-2-oxo-5-(N-cyclohexyl-N-methyl-
amino)-1H-1,4-benzo-diazepin-3-yl]-3-cyclohexylpropanamide

20

m.p. 170-171°C.

Anal. Calcd. for C₂₆H₃₈N₄O₂ 0.55 EtOAc 0.5 cyclohexane:

C, 70.81; H, 9.22; N, 10.59.

Found: C, 70.88; H, 9.2; N, 10.55%.

25

NMR (300 MHz, DMSO) δ: 8.46 (d, J = 8.1 Hz, 1H), 7.28-7.65 (m,
4H), 4.95 (d, J = 8.1 Hz, 1H), 3.1-3.4 (m, 1H), 3.26 (s, 3H), 2.64 (s,
3H), 2.17 (t, J = 7.1 Hz, 2H), 0.7-1.9 (m, 23H).

EXAMPLE 10

30

(±)-N-[2,3-Dihydro-1-methyl-2-oxo-5-(N,N-diethylamino)-1H-1,4-
benzodiazepin-3-yl]-5-phenylpentanamide

m.p. 134-135°C.

- 35 -

Anal. Calcd. for $C_{25}H_{32}N_4O_2 \cdot 0.25 H_2O$:

C, 70.64; H, 7.71; N, 13.18.

Found: C, 70.7; H, 7.94; N, 13.16%.

- 5 NMR (300 MHz, DMSO) δ : 8.48 (d, $J = 8.1$ Hz, 1H), 7.1-7.6 (m, 9H), 4.97 (d, $J = 8.1$ Hz, 1H), 3.25-3.4 (m, 2H), 3.27 (s, 3H), 2.95-3.1 (m, 2H), 2.53 (t, $J = 7.1$ Hz, 2H), 2.20 (t, $J = 7.1$ Hz, 2H), 1.4-1.6 (m, 4H), 0.97 (t, $J = 7.1$ Hz, 6H).

10

EXAMPLE 11

(\pm)-N-[2,3-Dihydro-1-methyl-2-oxo-5-(N,N-diethylamino)-1H-1,4-benzodiazepin-3-yl]-2-cyclohexylacetamide

- 15 m.p. 192-193°C.

Anal. Calcd. for $C_{22}H_{32}N_4O_2 \cdot 0.9 H_2O$:

C, 65.93; H, 8.5; N, 13.98.

Found: C, 65.91; H, 7.9; N, 14.05%.

- 20 NMR (300 MHz, DMSO) δ : 8.43 (d, $J = 8.1$ Hz, 1H), 7.2-7.65 (m, 4H), 4.96 (d, $J = 8.1$ Hz, 1H), 3.2-3.3 (m, 2H), 3.27 (s, 3H), 2.9-3.1 (m, 2H), 1.95-2.1 (m, 2H), 0.8-1.7 (m, 17H).

EXAMPLE 12

25

(\pm)-N-[2,3-Dihydro-1-methyl-2-oxo-5-(N,N-diethylamino)-1H-1,4-benzodiazepin-3-yl]-3-cyclohexylpropanamide

m.p. 209-210°C.

- 30 Anal. Calcd. for $C_{23}H_{34}N_4O_2$:

C, 69.31; H, 8.6; N, 14.06.

Found: C, 69.38; H, 8.69; N, 13.69%.

- 36 -

NMR (300 MHz, DMSO) δ : 8.45 (d, J = 8.1 Hz, 1H), 7.2-7.65 (m, 4H), 4.96 (d, J = 8.1 Hz, 1H), 3.2-3.4 (m, 2H), 3.27 (s, 3H), 2.9-3.1 (m, 2H), 2.17 (t, J = 7.3 Hz, 2H), 0.7-1.7 (m, 19H).

- 5 By using hexahydroazepine in place of N-methylcyclohexylamine, and the appropriate acid the following compounds were prepared using the processes substantially as described in Example 1.

EXAMPLE 13

10

N-[2,3-Dihydro-1-methyl-2-oxo-5-(hexahydroazepin-1-yl)-1
H-1,4-benzodiazepin-3-yl]-2-cyclohexyl acetamide

m.p. 205-207°C.

15

Anal. Calcd. for C₂₄H₃₄N₄O₂ · 0.25 H₂O

C, 70.21; H, 8.35; N, 13.65.

Found: C, 69.44; H, 8.38; N, 13.50%.

20

NMR (300 MHz, CD₃OD) δ 0.95-2.83 (m, 19H), 2.08 (m, 2H), 3.41 (m, 4H), 4.94 (s, 3H), 5.09 (d, 1H), 7.29-7.63 (m, 4H).

EXAMPLE 14

25

(+)-3-cyclohexyl-N-[2,3-Dihydro-1-methyl-2-oxo-5-(hexahydro-
azepin-1-yl)-1H-1,4-benzodiazepin-3-yl]propanamide

m.p. 100-102°C.

Anal. Calcd. for C₂₅H₃₆N₄O₂ · HCl, 0.50 H₂O; 0.30 EtOAc:

C, 63.38; H, 8.2; N, 11.28.

30

Found: C, 63.36; H, 8.09; N, 11.3%.

δ : 8.54 (d, J = 9.0 Hz, 1H), 7.79-7.74 (m, 1H), 7.54 (d, J = 7.8 Hz, 1H), 7.49-7.40 (m, 2H), 5.69 (dd, J = 9.0, 5.6 Hz, 2H), 4.55-4.4 (m, 1H), 4.20-4.00 (m, 1H), 3.54-3.49 (m, 2H), 3.46 (s, 3H), 2.47-2.41 (m, 2H),

- 37 -

2.09-1.97 (m, 5H), 1.82-1.51 (m, 10H), 1.28-1.12 (m, 4H), 0.93-0.85 (m, 2H).

By using N-methylbenzylamine in place of N-methylcyclohexylamine, and the appropriate acid, the following compounds were prepared using the processes substantially as described in Example 1.

EXAMPLE 15

N-[2,3-Dihydro-1-methyl-2-oxo-5-(N,N-dimethylamino)-1H-1,4-benzodiazepin-3-yl]-3-cyclohexylpropanamide

m.p. 197.5-198°C.

Anal. Calcd. for C₂₁H₃₀N₄O₂.0.05

C, 68.08; H, 8.16; N, 15.12.

Found: C, 67.91; H, 8.17; N, 14.95%.

NMR (300 MHz, CDCl₃) δ 0.81-1.77 (m, 13H), 2.30 (m, 2H), 2.86 (s, 6H), 3.21 (s, 3H), 5.26 (d, 1H), 6.99 (d, 1H), 7.22-7.52 (m, 4H)

2,3-Dihydro-1-benzyl-2-oxo-1H-1,4-benzodiazepine-2,5-dione was prepared from N-benzylisatoic anhydride (Transworld Chemicals) and glycine using the method described by Bock *et al.*, *J. Org. Chem.* 52:1644, (1987). This was then reacted with dimethylamine in place of the N-methyl cyclohexylamine, and the appropriate acid substantially as described above in Example 1 to give the following compounds:

EXAMPLE 16

N-[2,3-Dihydro-1-propyl-2-oxo-5-(N,N-dimethylamino)-1H-1,4-benzodiazepin-3-yl]-2-cyclohexylacetanide

- 38 -

m.p. 196-197°C.

Anal. Calcd. for C₂₂H₃₂N₄O₂:


C, 68.72; H, 8.39; N, 14.57.

Found: C, 68.54; H, 8.31; N, 14.44%.

5 NMR (300 MHz, DMSO) δ : 0.59-2.12 (m, 16H), 2.78 (m, 2H), 3.32 (s, 6H), 3.58 (m, 1H), 4.19 (m, 1H), 4.95 (d, 1H), 7.30-7.62 (m, 4H), 8.46 (d, 1H).

10 Additional compounds which can be used to selectively blockade the IKs channel also include for example:

EXAMPLE 17



25 (E)-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-phenyl-2-propenamide

A solution of (E)-3-phenyl-2-propenoyl chloride (367 mg, 2.2 mmol) in methylene chloride (1 mL) was added to a solution of 3(R)-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (*J. Org. Chem.* 1987, 52:3232-3239) (531 mg, 2.0 mmol) and triethylamine (307 μ L, 225 mg, 2.2 mmol) in methylene chloride (10 mL). The mixture was stirred at room temperature for 25 min. and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with CH₂Cl₂/Et₂O (95:5) and the residue was triturated with Et₂O. The

30

- 39 -

solid was collected and dried in vacuo at 70°C to give (E)-(+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-phenyl-2-propenamide as a colorless solid (170 mg, 21%), m.p. 140-142°C, $[\alpha]_D^{25} +86.7^\circ$ (c=0.173, CH₂Cl₂).

5 δ_H (CDCl₃) 7.70-7.26 (16H, m), 6.63 (1H, d, J 15.6 Hz), 5.68 (1H, d, J 8.3 Hz), and 3.50 (3H, s).

Anal. Calcd. for C₂₅H₂₁N₃O₂·0.15 (C₂H₅)₂O:

C, 75.63; H, 5.58; N, 10.33.

Found: C, 75.29; H, 5.57; N, 10.33%.


10

Employing the procedure substantially as described above, but substituting an appropriate acid chloride for the (E)-3-phenyl-2-propenoyl chloride, the following compounds were prepared:

15

EXAMPLE 18

20



25 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]benzamide

m.p. 224-225°C, $[\alpha]_D^{25} +89.2^\circ$ (c = 0.141, CH₂Cl₂).

δ_H (CDCl₃) 8.04 (1H, d, J 8.1 Hz), 7.96 (2H, d, J 6.8 Hz), 7.64-7.36 (10H, m), 7.27 (2H, t, J 7.6 Hz), 5.74 (1H, d, J 7.8 Hz), and 3.51 (3H, s).


30

Anal. Calcd. for C₂₃H₁₉N₃O₂·0.20H₂O:

C, 74.06; H, 5.24; N, 11.26.

Found: C, 74.13; H, 5.12; N, 11.16%.

- 40 -

EXAMPLE 19

First diastereoisomer to elute:

(-)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl](trans-2-phenyl-1-cyclopropane)carboxamide

m.p. 180-181°C, $[\alpha]_D -155.8^\circ$ (c = 0.434, CH₂Cl₂).

δ_H (CDCl₃) 7.62-7.09 (15H, m), 5.59 (1H, d, J 8.1 Hz), 3.47 (3H, s), 2.52-2.45 (1H, m), 1.90-1.84 (1H, m), 1.69-1.56 (1H, m), and 1.38-1.32 (1H, m).

Anal. Calcd. for C₂₆H₂₃N₃O₂·0.25H₂O:

C, 75.43; H, 5.72; N, 10.15.

Found: C, 75.38; H, 5.64; N, 9.94%.

Second diastereoisomer to elute:

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl](trans-2-phenyl-1-cyclopropane)carboxamide

m.p. 104-107°C, $[\alpha]_D +328.2^\circ$ (c = 0.098, CH₂Cl₂).

δ_H (CDCl₃) 7.62-7.13 (15H, m), 5.60 (1H, d, J 8.3 Hz), 3.48 (3H, s), 2.59-2.54 (1H, m), 1.93-1.87 (1H, m), 1.62-1.56 (1H, m, overlaps with water), and 1.33-1.25 (1H, m).

Anal. Calcd. for C₂₆H₂₃N₃O₂·0.50H₂O·0.45PhCH₃:

C, 76.13; H, 5.95; N, 9.14.

Found: C, 76.10; H, 5.94; N, 9.17%.

- 41 -

EXAMPLE 20


(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]-1H-indole-2-carboxamide

5 m.p. 167-177°C, $[\alpha]_D^{+113}$ (c = 1.103, CH₂Cl₂)
 δ_H (CDCl₃) 9.15 (1H, br s), 8.10 (1H, d, J 9.0 Hz), 7.75-7.10 (14H, m),
5.75 (1H, d, J 9.0 Hz), and 3.50 (3H, s).

Anal. Calcd. for C₂₅H₂₀N₄O₂:

C, 73.51; H, 4.94; N, 13.72.

10 Found: C, 73.31; H, 4.80; N, 13.62%.

EXAMPLE 21

20 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]heptanamide

m.p. 49-54°C, $[\alpha]_D^{+69.5}$ (c=1.000, MeOH).


Anal. Calcd. for C₂₃H₂₇N₃O₂·0.40H₂O:

25 C, 71.81; H, 7.28; N, 10.92.

Found: C, 71.90; H, 7.09; N, 10.85%.

30

- 42 -

EXAMPLE 22


10 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]hexanamide

[α]_D +72.6° (c=0.920, MeOH).

Anal. Calcd. for C₂₂H₂₅N₃O₂:

C, 72.70; H, 6.93; N, 11.56.

15 Found: C, 72.44; H, 6.75; N, 11.25%.

EXAMPLE 23

20
25 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]pentanamide


[α]_D +68.2° (c=1.310, MeOH).

Anal. Calcd. for C₂₁H₂₃N₃O₂·0.25CHCl₃:

30 C, 68.21; H, 6.26; N, 11.26.

Found: C, 68.2; H, 6.29; N, 11.17%.

- 43 -

EXAMPLE 24(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-phenylpropanamide

Oxalyl chloride (158 μ L, 230 mg, 1.81 mmol) was added to a mixture of 3-phenylpropanoic acid (249 mg, 1.66 mmol) and DMF (1 drop) in THF (10 mL) and the mixture was stirred at room temperature for 40 min. 3(R)-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (*J. Org. Chem.* 1987, 52:3232-3239) (400 mg, 1.51 mmol) and triethylamine (252 μ L, 183 mg, 1.81 mmol) were added and the mixture was stirred at room temperature for 18 h. The mixture was poured into saturated aqueous sodium hydrogen carbonate (20 mL) and extracted with ethyl acetate (3 x 20 mL). The combined organic fractions were dried (Na_2SO_4) and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$ (95:5) and the residue was recrystallized from toluene/hexane to give (+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-phenylpropanamide as a colorless solid (380 mg, 63%), m.p. 179°C, $[\alpha]_D^{+100.4^\circ}$ ($c = 0.225$, CH_2Cl_2).

δ_{H} (CDCl_3) 7.62-7.57 (2H, m), 7.47-7.21 (13H, m), 5.54 (1H, d, J 8.1 Hz), 3.47 (3H, s), 3.03 (2H, t, J 7.8 Hz), and 2.73-2.67 (2H, m).


Anal. Calcd. for $\text{C}_{25}\text{H}_{23}\text{N}_3\text{O}_2 \cdot 0.15\text{H}_2\text{O}$:

C, 75.04; H, 5.87; N, 10.50.

Found: C, 75.06; H, 5.78; N, 10.55%.

- 44 -

Employing the procedure substantially as described above, but substituting an appropriate carboxylic acid for the 3-phenylpropanoic acid, the following compounds were prepared:

EXAMPLE 25

E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(3,4-dichlorophenyl)-2-propenamide


m.p. 145-147°C, $[\alpha]_D^{25} +77.8^\circ$ (c=0.126, CH₂Cl₂).

δ_H (CDCl₃) 7.64-7.25 (14H, m), 6.61 (1H, d, J 15.6 Hz), 5.65 (1H, d, J 8.0 Hz), and 3.50 (3H, s).

Anal. Calcd. for C₂₅H₁₉N₃O₂Cl₂:

C, 64.67; H, 4.12; N, 9.05.

Found: C, 64.57; H, 4.25; N, 9.01%.

EXAMPLE 26

E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)-2-propenamide

- 45 -

m.p. 137-139°C, $[\alpha]_D^{+66.0^\circ}$ ($c=0.144$, CH_2Cl_2).


δ_H (CDCl_3) 8.02 (1H, d, J 15.6 Hz), 7.73-7.26 (13H, m), 6.66 (1H, d, J 15.6 Hz), 5.81 (1H, d, J 8.8 Hz), and 3.53 (3H, s).

Anal. Calcd. for $\text{C}_{25}\text{H}_{19}\text{Cl}_2\text{N}_3\text{O}_2$:

5 C, 64.67; H, 4.12; N, 9.05.

Found: C, 64.28; H, 4.24; N, 8.83%.

EXAMPLE 27



E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(4-methylphenyl)-2-propenamide

m.p. 133-135°C, $[\alpha]_D^{+90.4^\circ}$ ($c=0.125$, CH_2Cl_2).

20 δ_H (CDCl_3) 7.68-7.19 (15H, m), 6.59 (1H, d, J 15.6 Hz), 5.70 (1H, d, J 8.0 Hz), 3.50 (3H, s), and 2.38 (3H, s).


Anal. Calcd. for $\text{C}_{26}\text{H}_{23}\text{N}_3\text{O}_2$:

C, 76.26; H, 5.66; N, 10.26.

Found: C, 75.93; H, 5.82; N, 10.10%.

25

EXAMPLE 28



- 46 -

E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(4-methoxyphenyl)-2-propenamide

m.p. 129-133°C, $[\alpha]_D^{+89.9^\circ}$ (c 0.188, CH₂Cl₂).


δ_H (CDCl₃) 7.65-7.24 (14H, m), 6.92 (1H, d, J 8.8 Hz), 6.50 (1H, d, J 15.6 Hz), 5.69 (1H, d, J 8.0 Hz), 3.84 (3H, s), and 3.50 (3H, s)..

Anal. Calcd. for C₂₆H₂₃N₃O₃·0.30H₂O:

C, 72.48; H, 5.52; N, 9.75.

Found: C, 72.75; H, 5.60; N, 9.36%.

EXAMPLE 29



(+)-5-Chloro-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]indole-2-carboxamide

m.p. 160-164°C, $[\alpha]_D^{+103.8^\circ}$ (c = 0.160, CH₂Cl₂).


δ_H (CDCl₃) 9.71 (1H, br s), 8.13 (1H, d, J 7.8 Hz), 7.68-7.09 (13H, m), 5.75 (1H, d, J 7.8 Hz), and 3.53 (3H, s).

Anal. Calcd. for C₂₅H₁₉ClN₄O₂·0.25H₂O·0.15PhCH₃:

C, 67.84; H, 4.49; N, 12.15.

Found: C, 67.80; H, 4.41; N, 12.07%.

- 47 -


EXAMPLE 30

10 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]-2,2-diphenylethanamide
m.p. 200-201°C, $[\alpha]_D^{+97.0^\circ}$ (c = 0.168, CH₂Cl₂).
 δ_H (CDCl₃) 7.60-7.22 (20H, m), 5.58 (1H, d, J 8.1 Hz), 5.08 (1H, s),
and 3.44 (3H, s).

15 Anal. Calcd. for C₃₀H₂₅N₃O₂·0.15PhCH₃:

C, 78.79; H, 5.55; N, 8.88.

Found: C, 78.81; H, 5.63; N, 9.07%.

EXAMPLE 31


20 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]-2-phenylethanamide
25 m.p. 241-242°C (dec.), $[\alpha]_D^{+85.5^\circ}$ (c = 0.159, CH₂Cl₂).
 δ_H (CDCl₃) 7.59-7.55 (3H, m), 7.46-7.22 (12H, m), 5.51 (1H, d, J
8.1Hz), 3.72 (2H, s), and 3.44 (3H, s).

- 48 -

Anal. Calcd. for $C_{24}H_{21}N_3O_2 \cdot 0.55H_2O$:

C, 73.28; H, 5.66; N, 10.68.

Found: C, 73.25; H, 5.38; N, 10.47%.


EXAMPLE 32

(+)-3-Cyclohexyl-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide

m.p. 144.5-145.5°C, $[\alpha]_D^{+83.1^\circ}$ (c = 0.116, CH_2Cl_2). δ_H ($CDCl_3$) 7.62-7.56 (3H, m), 7.46-7.21 (7H, m), 5.55 (1H, d, J 8.3 Hz), 3.48 (3H, s), 2.41-2.36 (2H, m), 1.77-1.58 (7H, m), 1.31-1.16 (4H, m), and 0.98-0.90 (2H, m).Anal. Calcd. for $C_{25}H_{29}N_3O_2$:

C, 74.41; H, 7.24; N, 10.41.

Found: C, 74.46; H, 7.27; N, 10.58%.

EXAMPLE 33

- 49 -

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(4-cyanophenyl)propanamide

m.p. 81-85°C, $[\alpha]_D^{+91.0^\circ}$ (c = 0.111, CH₂Cl₂).


δ_H (CDCl₃) 7.64-7.55 (4H, m), 7.48-7.16 (10H, m), 5.50 (1H, d, J 8.3 Hz), 3.47 (3H, s), 3.08 (2H, t, J 7.6 Hz), and 2.74-2.69 (2H, m).

Anal. Calcd. for C₂₆H₂₂N₄O₂·0.60H₂O·0.50PhCH₃:

C, 73.93; H, 5.62; N, 11.69.

Found: C, 73.98; H, 5.61; N, 11.71%.

EXAMPLE 34



(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(3-chlorophenyl)propanamide

m.p. 157-159°C, $[\alpha]_D^{+90.7^\circ}$ (c = 0.134, CH₂Cl₂).


δ_H (CDCl₃) 7.62-7.57 (3H, m), 7.47-7.12 (11H, m), 5.53 (1H, d, J 8.1 Hz), 3.47 (3H, s), 3.00 (2H, t, J 7.3 Hz), and 2.71-2.66 (2H, m).

Anal. Calcd. for C₂₅H₂₂ClN₃O₂·0.55H₂O:

C, 67.96; H, 5.27; N, 9.51.

Found: C, 67.99; H, 5.18; N, 9.26%.

- 50 -

EXAMPLE 35

E-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(3-bromophenyl)-2-propenamide


m.p. 221-223 d°C, $[\alpha]_D^{+65.5^\circ}$ (c = 0.206, CH₂Cl₂).

δ_H (CDCl₃) 7.69 (1H, br s), 7.64-7.57 (4H, m), 7.51-7.37 (6H, m), 7.29-7.19 (4H, m), 6.62 (1H, d, J 15.6 Hz), 5.66 (1H, d, J 8.1 Hz), and 3.50 (3H, s).

Anal. Calcd. for C₂₅H₂₀BrN₃O₂·0.35H₂O·0.20PhCH₃:

C, 63.54; H, 4.46; N, 8.42.

Found: C, 63.50; H, 4.39; N, 8.42%.

EXAMPLE 36

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-4-phenylbutanamide

m.p. 65-74°C, $[\alpha]_D^{+77.4^\circ}$ (c = 0.155, CH₂Cl₂).

- 51 -

δ_H (CDCl₃) 7.62-7.56 (3H, m), 7.46-7.19 (12H, m), 5.55 (1H, d, J 8.1 Hz), 3.47 (3H, s), 2.71 (2H, t, J 7.6 Hz), 2.42-2.37 (2H, m), and 2.09-2.01 (2H, m).

Anal. Calcd. for C₂₆H₂₅N₃O₂·0.30H₂O:


5 C, 74.91; H, 6.19; N, 10.08.

Found: C, 74.93; H, 6.05; N, 10.07%.

EXAMPLE 37

10

15



(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]-3-(3-cyanophenyl)propanamide

m.p. 110-112°C, [α]_D +84.2° (c = 0.202, CH₂Cl₂).

20 δ_H (CDCl₃) 7.63-7.22 (14H, m), 5.51 (1H, d, J 8.1 Hz), 3.47 (3H, s),
3.06 (2H, t, J 7.8 Hz), and 2.74-2.68 (2H, m).

Anal. Calcd. for C₂₆H₂₂N₄O₂·0.50H₂O:


C, 72.37; H, 5.37; N, 12.98.

Found: C, 72.52; H, 5.12; N, 12.59%.

25

EXAMPLE 38

30



- 52 -

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-4-methylpentanamide

m.p. 123-125°C, $[\alpha]_D^{25} +66.8^\circ$ (c=0.500, MeOH).


Anal. Calcd. for C₂₂H₂₅N₃O₂·0.45H₂O:

5 C, 71.12; H, 7.03; N, 11.31.

Found: C, 71.08; H, 6.81; N, 11.42%.

EXAMPLE 39

10



(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2,3-dihydrobenzofuran-2-carboxamide

Diisopropylethylamine (0.3 mL, 223 mg, 1.72 mmol) was added to a stirred, cooled (0°C) solution of 3(R)-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (*J. Org. Chem.* 1987, 52:3232-3239) (400 mg, 1.5 mmol), 2,3-dihydrobenzofuran-2-carboxylic acid (274 mg, 1.7 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodi-imide hydrochloride (583 mg, 3.0 mmol), and 1-hydroxybenzotriazole (479 mg, 3.1 mmol) in DMF (4.5 mL). The mixture was stirred at room temperature for 18 h., poured into aqueous hydrochloric acid (3M, 12 mL) and extracted with ethyl acetate (3 x 20 mL). The combined organic fractions were washed with saturated aqueous sodium hydrogen carbonate (20 mL) and brine (20 mL), dried (MgSO₄) and evaporated under reduced pressure. The residue was crystallized from 2-chloro-2-methyl-propane/hexane to give (+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2,3-dihydrobenzofuran-2-carboxamide as a colorless solid (156 mg, 25%), m.p. 141-180°C, $[\alpha]_D^{25} +127.1^\circ$ (c=0.425, CHCl₃).

20

25

30

- 53 -

δ H (CDCl₃) (3:1 Mixture of diastereoisomers) 8.44 (1H, m), 7.65-6.91 (13H, m), 5.52 (1H, m), 5.28 (1H, m), and 3.70-3.40 (5H, m).

Anal. Calcd. for C₂₅H₂₁N₃O₃·0.25 Hexane

C, 73.50; H, 5.70; N, 9.71.

5 Found: C, 74.12; H, 5.57; N, 9.71%.

EXAMPLE 40

Although not a pure IKs inhibitor, the synthesis of (+)-N-
10 [(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-diazepin-3-yl]-1'-(1,1-dimethylethoxycarbonyl)spiro(cyclohexan-4,4'-piperidine)-1-carboxamide illustrates a procedure for preparing other compounds which are active.

15 Step A:



25 Diethyl 1-benzylpiperidine-4,4-diacetate


Ethanol (120 mL) was cooled in ice and ammonia bubbled through to give a saturated solution. 1-Benzyl-4-piperidone (40.0 g, 211 mmol) and ethyl cyanoacetate (47.8 g, 423 mmol) were added, the reaction vessel stoppered and stored at 0°C overnight. The solid was
30 collected, washed with ethanol and ether and dried *in vacuo* to give a yellow solid (68.86 g). The solid (58.86 g) was dissolved in a mixture of sulfuric acid (70 mL, 98%) and water (60 mL) and heated under reflux for three days the mixture cooled and most of the water evaporated. The residue was azeotroped with ethanol (4x750 mL),

- 54 -

further ethanol (500 mL) added and the mixture heated under reflux for 20h, cooled in ice and sodium carbonate (100 g) added slowly with vigorous stirring. The ethanol was evaporated under reduced pressure, water (800 mL) added and the mixture extracted with methylene chloride (3x400 mL). The combined organic extracts were dried (Na₂SO₄) and the solvent evaporated to give Diethyl 1-benzylpiperidine-4,4-diacetate (37.51 g). A small portion of this was purified by flash column chromatography.

NMR (300 MHz, CDCl₃) δ : 7.2-7.4 (m, 5H), 4.11 (q, J=7.3Hz, 4H), 3.50 (s, 2H), 2.56 (s, 4H), 2.4 (m, 4H), 1.7 (m, 4H), 1.24 (t, J=7.3Hz, 6H).

Step B:



20 1-Benzylpiperidine-4,4-diethanol


A solution of the diester (12.2 g, 35 mmol) in ether (25 mL) was added to a cooled (-30°C) and stirred suspension of LiAlH₄ (2.1 g, 55 mmol) in ether (400 mL), under argon. THF (60 mL) was added and the reaction mixture allowed to warm to room temperature. After recooling to 0°C, water (2.2 mL), 1M NaOH (4.4 mL) and water (5 mL) were added, the reaction mixture stirred vigorously for 30 min and the solid filtered off, washing well with ether. The combined filtrates were evaporated to afford a white solid which was triturated with ether to give 8 g of 1-benzylpiperidine-4,4-diethanol.

30

m.p. 75-78°C


NMR (300 MHz, CDCl₃) δ : 7.2-7.4 (m, 5H), 3.7 (t, J = 6.8 Hz, 4H), 3.52 (s, 2H), 2.7 (brs, 2H), 2.43 (m, 4H), 1.66 (t, J = 6.8 Hz, 4H), 1.5 (m, 4H).

- 55 -

Step C:1-t-Butoxycarbonylpiperidine-4,4-diethanol

The benzylamine (2.07 g, 7.9 mmol) was dissolved in methanol (60 mL), BOC₂O (1.72 g, 7.9 mmol) added and the mixture hydrogenated at 50 psi over 10% palladium hydroxide on charcoal (200 mg) for 18 hours. The reaction mixture was filtered through celite, washed with methanol and the filtrate evaporated to give 1-t-Butoxycarbonylpiperidine-4,4-diethanol (2.0 g).

NMR (300 MHz, CDCl₃) δ : 3.7 (m, 4H), d 3.3 (m, 6H), 1.65 (t, J = 6.8 Hz, 4H), 1.41 (s, 9H).


Step D:1-t-Butoxycarbonylpiperidine-4,4-diethanol, bis(methanesulfonate)

The diol (2.41 g, 8.9 mmol) was dissolved in dichloromethane (50 mL), the solution cooled to -20°C under argon before addition of triethylamine (3.7 mL, 26 mmol) and methanesulfonyl chloride (1.6 mL, 20 mmol). After 30 min., the reaction mixture was poured into ice cold 10% citric acid and extracted with ether (X3). The combined extracts were washed with water, saturated NaHCO₃ and brine, dried (MgSO₄) and the solvent evaporated to afford 1-t-Butoxycarbonylpiperidine-4,4-diethanol, bis(methanesulfonate) (3.2 g).

- 56 -

NMR (300 MHz, CDCl₃) δ : 4.32 (t, J = 7.1 Hz, 4H), 3.4 (m, 4H), 3.04 (s, 6H), 1.89 (t, J = 7.1 Hz, 4H).

5 Step E:




15 Diethyl 3-t-butyloxycarbonyl-3-azaspiro[5.5]undecane-9,9-dicarboxylate

To a slurry of 60% NaH (2.04 g, 0.51 mole) in toluene (160 mL), under argon, was slowly added diethyl malonate (3.72 mL, 24.3 mmol). The mixture was cooled to 0°C and the bis-mesylate 1 (7.0 g, 16.3 mmol) added as a solid and the mixture heated to reflux for 18 hours. The reaction was quenched into 10% citric acid (100 mL) and the product extracted with CH₂Cl₂ (2x150 mL). The extracts were dried (Na₂SO₄), concentrated to an oil, and chromatographed on silica to give 3.83 g (60%) of diethyl 3-t-butyloxycarbonyl-3-azaspiro[5.5]undecane-9,9-dicarboxylate.

25 ¹H NMR (CDCl₃) δ : 1.22 (t, 6H), 1.4 (s, 9H), 2.0 (m, 4H), 3.35 (m, 4H), 4.2 (q, 4H).

30


- 57 -

Step F:

10 3-t-Butyloxycarbonyl-3-azaspiro[5.5]undecane-9-carboxylic acid

To a solution of the diester 2 (3.69 g, 0.0093 m) in THF (50 mL) was added 1N LiOH (47 mL). The reaction was stirred for 3 days at 25°C, diluted with water (50 mL) and pH adjusted to 2.2 with KHSO₄. The product was extracted into ethyl acetate (2x75 mL), dried (Na₂SO₄), and concentrated to a foam (3.5 g). The solid was melted in a flask at 140°C for 2 hours, cooled and the oil dissolved in THF (15 mL), 1N LiOH (10 mL) added and mixture stirred overnight at 30°C. The reaction was concentrated to remove THF, diluted with water (20 mL) and washed with diethyl ether (10 mL). The pH was adjusted to 2.5 with KHSO₄ and product extracted (3x50 mL) with ethyl acetate. The extracts were dried (Na₂SO₄), filtered and concentrated to yield 3-t-Butyloxycarbonyl-3-azaspiro[5.5]undecane-9-carboxylic acid as a foam (2.48 g, 90%).

25 ¹H NMR (CDCl₃, partial) δ: 1.45 (s, 9H), 3.4 (m, 4H).

Step G:

- 58 -

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-1'-(1,1-dimethylethoxycarbonyl)spiro(cyclohexan-4,4'-piperidine)-1-carboxamide

m.p. 135-138°C, $[\alpha]_D^{25} +58.8^\circ$ (C=0.925, CHCl₃).

5 δ_H (CDCl₃) 7.61-7.23 (10H, m), 5.54 (1H, d, J 9.0 Hz), 3.47 (3H, s), 3.37 (4H, m), 2.28 (1H, m), and 1.81-1.18 (21H, s).

Anal. Calcd. for C₃₂H₄₀N₄O₄:

C, 70.56; H, 7.40; N, 10.29.

Found: C, 70.21; H, 7.40; N, 10.16%.


10

Employing the procedure substantially as described in Example 43 but substituting an appropriate acid for the 2,3-dihydro-benzofuran-2-carboxylic acid, the following compounds were prepared:

15

EXAMPLE 41

20



25 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(furan-2-yl)propanamide

m.p. 115-118°C, $[\alpha]_D^{25} +65.8^\circ$ (c=0.800, CHCl₃).


δ_H (CDCl₃) 7.62-7.26 (11H, m), 6.28 (1H, dd, J 3.2, 2.0 Hz), 6.08 (1H, dd, J 3.2, 0.7 Hz), 5.58 (1H, d, J 8.1 Hz), 3.48 (3H, s), 3.04 (2H, t, J 7.6 Hz), and 2.75 (2H, m).

30 Anal. Calcd. for C₂₃H₂₁N₃O₃·0.3Hexane:

C, 72.07; H, 6.15; N, 10.17.

Found: C, 71.78; H, 6.30; N, 9.77%.

- 59 -

EXAMPLE 42

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]cyclohexylcarboxamide

m.p. 213-214°C, $[\alpha]_D^{25} +62.4^\circ$ (c=1.000, MeOH).

Anal. Calcd. for C₂₃H₂₄N₃O₂:


C, 73.77; H, 6.46; N, 11.22.

Found: C, 73.86; H, 6.81; N, 11.15%.

EXAMPLE 43

L-742192-000C001

OC1568-5113



(E)-(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]-3-(3,4-methylenedioxyphenyl)-2-propenamide

m.p. 143-145°C, $[\alpha]_D^{25} +62.3^\circ$ (c=0.960, MeOH).

Anal. Calcd. for C₂₅H₂₁N₃O₄·0.10H₂O·0.20Et₂O:


C, 69.78; H, 5.27; N, 9.46.

Found: C, 69.78; H, 4.98; N, 9.28%.

- 60 -

EXAMPLE 44

(+)-N-[(3R)-2,3-Dihydro-2-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-(phenylamino)acetamide


Step A:

N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-bromoacetamide

Bromoacetyl bromide (165 μ L, 383 mg, 1.9 mmol) was added to an ice cooled solution of 3(R)-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (*J. Org. Chem.* 1987, 52:3232-3239) (500 mg, 1.88 mmol) and triethylamine (264 μ L, 192 mg, 1.9 mmol) in methylene chloride (10 mL) and the mixture was stirred at room temperature for 1 h. The mixture was washed with water (3 x 10 mL), dried (MgSO_4) and the solvent was evaporated under reduced pressure to give N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-bromoacetamide as a colorless foam (760 mg, 100%).

δ_{H} (CDCl_3) 8.24 (1H, d, J 7.8 Hz), 7.64-7.24 (9H, m), 5.48 (1H, d, J 7.8 Hz), 4.00 (2H, m), and 3.50 (3H, s).

- 61 -

Step B:

5
10 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-(phenylamino)acetamide

Aniline (297 μ L, 304 mg, 3.26 mmol) was added to a solution of N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-bromoacetamide (600 mg, 1.55 mmol) in ethanol (25 mL) and the mixture was heated under reflux for 24 h. The mixture was cooled and the solid was collected and recrystallized from ethanol (20 mL) to give (+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-(phenylamino)acetamide as a colorless solid (500 mg, 81%), m.p. 245-246°C, $[\alpha]_D +119^\circ$ (C=0.850, CHCl₃).

15
20 δ_H (CDCl₃) 8.26 (1H, d, J 8.3 Hz), 7.63-7.20 (12H, m), 6.81 (1H, t, J 7.3 Hz), 6.72 (2H, d, J 7.6 Hz), 5.56 (1H, d, J 8.3 Hz), 3.95 (2H, d, J 1.5 Hz), and 3.45 (3H, s).

Anal. Calcd. for C₂₄H₂₂N₄O₂:


25 C, 72.34; H, 5.57; N, 14.06.

Found: C, 72.37; H, 5.59; N, 14.32%.


Employing the procedure substantially as described above, but substituting 2-chloroaniline or 4-(trifluoromethyl)aniline for the aniline, the following compounds were prepared:

30

- 62 -

EXAMPLE 45

- 10 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-
diazepin-3-yl]-2-(2-chlorophenylamino)acetamide
m.p. 222-224°C, $[\alpha]_D^{+111}$ (c=0.973, CHCl₃).
 δ_H (CDCl₃) 8.15 (1H, d, J 8.3 Hz), 7.60-7.16 (12H, m), 6.71 (2H, m),
5.57 (1H, d, J 8.3 Hz), 4.01 (2H, d, J 2.7 Hz), and 3.45 (3H, s).
15 Anal. Calcd. for C₂₄H₂₁ClN₄O₂:
C, 66.59; H, 4.89; N, 12.94.
Found: C, 66.40; H, 4.94; N, 12.92%.

EXAMPLE 46

- 20
25
30 (+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-
benzo-diazepin-3-yl]-2-(phenoxy)acetamide
Phenol (104 mg, 1.1 mmol) was added to a suspension of
sodium hydride (60% dispersion in mineral oil, 44 mg, 1.1 mmol) in
toluene (10 mL). When hydrogen evolution had stopped, N-[(3R)-2,3-
dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-
bromoacetamide (400 mg, 1.04 mmol) was added and the mixture was

- 63 -

stirred at room temperature for 18 h. The mixture was washed with water (3 x 15 mL), dried (MgSO₄) and the solvent was evaporated under reduced pressure. The residue was triturated with 2-propanol and the solid was collected and recrystallized from 2-propanol (5 mL) to give (+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-(phenoxy)acetamide as a colorless solid (112 mg, 27%), m.p. 126-128°C, [α]_D +81.6 (C=0.692, CHCl₃). δ _H (CDCl₃) 8.49 (1H, d, J 8.2 Hz), 7.64-7.01 (14H, m), 5.61 (1H, d, J 8.2 Hz), 4.65 (1H, d, J 14.6 Hz), 4.58 (1H, d, J 14.6 Hz), and 3.50 (3H, s).


Anal. Calcd. for C₂₄H₂₁N₃O₃:

C, 72.17; H, 5.30; N, 10.52.

Found: C, 71.84; H, 5.25; N, 10.41%.

Employing the procedure substantially as described above, but substituting 2,4-dichlorophenol, thiophenol or 2,4-dichlorothiophenol for the phenol, the following compounds were prepared:

EXAMPLE 47



(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-2-(phenylthio)acetamide


[α]_D +104.9° (c=0.316, CHCl₃).

δ _H (CDCl₃) 8.50 (1H, d, J 9.0 Hz), 7.60-7.20 (14H, m), 5.50 (1H, d, J 9.0 Hz), 3.75 (2H, m), and 3.45 (3H, s).

Anal. Calcd. for C₂₄H₂₁N₃O₂S:

- 64 -

Found: C, 69.37; H, 5.10; N, 10.11.
C, 68.98; H, 5.06; N, 9.76%.

EXAMPLE 48

(+)-N-[(3R)-2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzo-diazepin-3-yl]-3-(phenylamino)propanamide


3-Bromopropionyl chloride (2.01 mL, 3.428 g, 20 mmol) was added to an ice cooled solution of 3(R)-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (*J. Org. Chem.* 1987, 52:3232-3239) (5.0 g, 18.8 mmol) and triethylamine (2.79 mL, 2.02 mg, 20 mmol) in methylene chloride (85mL) and the mixture was stirred at room temperature for 18 h. The mixture was washed with saturated aqueous sodium hydrogen carbonate (85 mL), water (2 x 85 mL), and brine (85 mL), dried (MgSO₄) and the solvent was evaporated under reduced pressure. A sample (0.5 g, 1.25 mmol) was dissolved in ethanol (25 mL), aniline (230 μ L, 233 mg, 2.5 mmol) was added and the mixture was heated under reflux for 70 h. The mixture was cooled and the solid was collected and recrystallized from ethanol to give (+)-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(phenylamino)propanamide as a colorless solid, m.p. 218-221°C, [α]_D +58.2° (c=0.585, CHCl₃). δ _H (CDCl₃) 7.60-6.71 (16H, m), 5.54 (1H, d, J 8.1 Hz), 3.54 (2H, t, J 6.1 Hz), 3.52 (3H, s), and 2.70 (2H, m).

- 65 -

Anal. Calcd. for $C_{25}H_{24}N_4O_2 \cdot 0.5EtOH$:

C, 71.70; H, 6.25; N, 12.87.

Found: C, 71.42; H, 5.98; N, 12.84%.

EXAMPLE 49

15 (-)-3-Cyclohexyl-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-4-oxido-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide

3-Chloroperoxybenzoic acid (80%, 0.32 g, 1.5 mmol) was added to a solution of (+)-3-cyclohexyl-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide (0.60 g, 1.5 mmol) in dichloromethane (25 mL) and the mixture was stirred at room temperature for 18 h. Further 3-chloroperoxybenzoic acid (80%, 0.1 g, 0.5 mmol) was added and the mixture was stirred for 24 h. The mixture was washed with saturated aqueous sodium hydrogen carbonate (4 x 25 mL), water (2 x 25 mL) and brine (25 mL), dried (MgSO₄) and the solvent was evaporated under reduced pressure. The residue was recrystallized from toluene/hexane (65:35) to give (-)-3-cyclohexyl-N-[(3R)-2,3-dihydro-1-methyl-2-oxo-4-oxido-5-phenyl-1H-1,4-benzodiazepin-3-yl]propanamide as colorless prisms, m.p. 222-224°C, $[\alpha]_D -80.7^\circ$ (c=1.15, CHCl₃).


30 δ_H (CDCl₃) 7.71-7.23 (10H, m), 6.01 (1H, d, J 9.3 Hz), 3.54 (3H, s), 2.48 (2H, m), and 1.76-0.89 (13H, m).

Anal. Calcd. for $C_{25}H_{29}N_3O_3 \cdot 0.5H_2O$:

C, 70.06; H, 7.06; N, 9.81.

Found: C, 70.10; H, 6.80; N, 9.79%.

- 66 -

EXAMPLE 50

10 (+)-Phenylmethyl N-[(3R)-2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl]carbamate

A mixture of (+)-phenylmethyl N-[(3R)-2,3-dihydro-1-methyl-5-phenyl-2-oxo-1H-1,4-benzodiazepin-3-yl]carbamate (4.0 g, 10 mmol) and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (4.5 g, 11 mmol) in toluene (100 mL) was heated under reflux for 75 min. The mixture was cooled and the volume was reduced to 30 mL by evaporation under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with EtOAc/Hexane (75:25) to give (+)-phenylmethyl N-[(3R)-2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl]carbamate as a solid, m.p. 128-131°C, $[\alpha]_D^{25} +22.5^\circ$ ($c=0.656$, CHCl_3). δ_H (CDCl_3) 7.65-7.26 (15H, m), 5.50 (1H, d, J 8.8 Hz), 5.14 (2H, s), and 3.86 (3H, s).


Anal. Calcd. for $\text{C}_{24}\text{H}_{21}\text{N}_3\text{O}_2\text{S} \cdot 0.25\text{H}_2\text{O}$:

25 C, 68.63; H, 5.16; N, 10.01.

Found: C, 68.28; H, 5.21; N, 10.06%.

30

- 67 -

EXAMPLE 51

10 3-Cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl)propanamide

Hydrogen bromide was bubbled at room temperature through a solution of (+)-phenylmethyl N-[(3R)-2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl]carbamate (0.9 g, 2.1 mmol), acetic acid (5 mL) and dichloromethane (5 mL). After 2 h., the solvent was evaporated under reduced pressure, ether was added and the solid was collected and dried *in vacuo*. A sample (0.58 g, 1.8 mmol) was suspended in THF (10 mL), triethylamine (0.24 mL, 1.8 mmol) was added and the mixture was stirred at room temperature for 3 h. In a separate flask, oxalyl chloride (0.20 mL, 0.29 g, 2.3 mmol) was added to a solution of cyclohexanepropionic acid (0.33 mL, 0.30 g, 1.9 mmol) and DMF (1 drop) in THF (10 mL) and the mixture was stirred at room temperature for 3 h. The two mixtures were combined, triethylamine (0.32 mL, 0.23 g, 2.3 mmol) was added and the mixture was stirred at room temperature for 2.5 h. The solvent was evaporated under reduced pressure, water was added and the mixture was extracted with ethyl acetate. The combined organic fractions were washed with water, saturated aqueous sodium hydrogen carbonate, water (2 x) and brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with CH₂Cl₂/MeOH (99.5:0.5) and the residue was recrystallized from EtOAc/Hexane to give 3-cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl)propanamide as a solid, m.p. 219-221°C.

- 68 -

δ H (CDCl₃) 7.95 (1H, br d, J 8.6 Hz), 7.65-7.30 (9H, m), 5.72 (1H, d, J 8.6 Hz), 3.87 (3H, s), 2.41 (2H, t, J 7.6 Hz), and 1.80-0.85 (13H, m).

Anal. Calcd. for C₂₅H₂₉N₃OS.0.25H₂O:


C, 70.81; H, 7.01; N, 9.91.

5 Found: C, 70.80; H, 6.91; N, 9.95%.

EXAMPLE 52

10

15



(E)- and (Z)-3-Cyclohexyl-N-(2,3-dihydro-2-hydroxyimino-5-phenyl-1H-1,4-benzodiazepin-3-yl)propanamide

20

A mixture of 3-cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl)propanamide (740 mg, 1.83 mmol), hydroxylamine hydrochloride (140 mg, 2 mmol) and triethylamine (280 μ L, 203 mg, 2 mmol) in methanol (15 mL)/THF (15 mL) was stirred at room temperature for 3 h. The solvent was

25 evaporated under reduced pressure and the residue was purified by flash column chromatography on silica gel, eluting with CH₂Cl₂/MeOH (98:2). The residue recrystallized from ethyl acetate. The first isomer to crystallize was recrystallized from ethyl acetate to give (E)-3-cyclohexyl-N-(2,3-dihydro-2-hydroxyimino-5-phenyl-1H-1,4-

30 benzodiazepin-3-yl)propanamide as a solid, m.p. 196°C.

δ H (d₆-DMSO) 12.20 (1H, s), 9.00 (1H, d, J 8.0 Hz), 7.70-7.30 (10H, m), 5.45 (1H, d, J 8.0 Hz), 2.30 (2H, m), and 1.80-0.75 (13H, m).

- 69 -


The second isomer to crystallize was recrystallized from methanol to give (Z)-3-cyclohexyl-N-(2,3-dihydro-2-hydroxyimino-5-phenyl-1H-1,4-benzodiazepin-3-yl)propanamide as a solid, m.p. 219°C. δ_H (d_6 -DMSO) 9.95 (1H, s), 8.95 (1H, s), 8.75 (1H, d, J 8.0 Hz), 7.50-7.00 (9H, m), 5.70 (1H, d, J 8.0 Hz), 2.25 (2H, m), and 1.75-0.75 (13H, m).

Anal. Calcd. for $C_{24}H_{28}N_4O_2$:

C, 71.26; H, 6.98; N, 13.85.

Found: C, 70.89; H, 6.99; N, 13.55%.

EXAMPLE 53



3-Cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-3-yl) propanamide

Freshly prepared Raney nickel (400 mg) was added to a solution of 3-cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-2-thioxo-1H-1,4-benzodiazepin-3-yl)propanamide (200 mg, 0.5 mmol) in ethanol (20 mL) and the mixture was stirred at room temperature for 2 h. The mixture was filtered and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with CH_2Cl_2 /MeOH (99.75:0.25) to give 3-cyclohexyl-N-(2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepin-3-yl) propanamide as a foam.

δ_H ($CDCl_3$) 7.60-6.80 (9H, m), 6.37 (1H, br d, J 6.6 Hz), 5.53 (1H, m), 3.60 (2H, m), 2.77 (3H, s), 2.21 (2H, t, J 8.0 Hz), and 1.85-0.80 (13H, m).


Anal. Calcd. for $C_{25}H_{31}N_3O \cdot 0.2CH_2Cl_2$:

- 70 -

Found: C, 74.45; H, 7.79; N, 10.34.
C, 74.68; H, 7.87; N, 10.23%.

EXAMPLE 54

1-(2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-thieno-[2,3-e]-1,4-diazepin-3-yl)-3-(3-methyl-phenyl)urea

Step A:(2-Amino-3-thienyl)phenylmethanone

Triethylamine (6.8 mL, 4.94 g, 49 mmol) was added to a heated (33°C) mixture of β -oxobenzene propanenitrile (18.6 g, 128 mmol) and 1,2-dithiane-2,5-diol (9.8 g, 64 mmol) in ethanol (120 mL) and the mixture was stirred at 50°C for 18 h. The mixture was cooled and the solvent was evaporated under reduced pressure.

Dichloromethane was added, the mixture was washed with aqueous hydrochloric acid (0.5M), aqueous sodium hydroxide (1M) and brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure.

The residue was recrystallized from acetonitrile (150 mL) to give (2-amino-3-thienyl)-phenylmethanone as an orange solid (5.7 g, 44%).
 δ_H (CDCl₃) 7.70-7.35 (5H, m), 6.95 (2H, br s), 6.90 (1H, d, J 6.3 Hz), and 6.15 (1H, d, J 6.3 Hz).


- 71 -

Step B:2,3-Dihydro-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one

A solution of 1,3-dihydro-1,3-dioxo-2H-isoindole-2-acetyl
5 chloride (8.6 g, 38 mmol) in dichloromethane (20 mL) was added
slowly to a cooled (0°C) mixture of (2-amino-3-thienyl)phenyl-
methanone (6.8 g, 33 mmol), pyridine (6.34 mL, 6.20 g, 78 mmol) and
4-dimethylamino-pyridine (0.79 g, 6.5 mmol) in dichloromethane (130
10 mL). The mixture was stirred at 0°C for 30 min., diluted with
dichloromethane (80 mL) and washed with aqueous hydrochloric acid
(1M), saturated aqueous sodium hydrogen carbonate and brine. The
mixture was dried (Na₂SO₄) and the solvent was evaporated under
reduced pressure. The residue was triturated with ethanol and the solid
15 was collected and dried in vacuo to give N-(3-benzoylthien-2-yl)-1,3-
dihydro-1,3-dioxo-2H-isoindole-2-acetamide as a solid (9.8 g, 76%).

A mixture of N-(3-benzoylthien-2-yl)-1,3-dihydro-1,3-
dioxo-2H-isoindole-2-acetamide (10.9 g, 28 mmol) and hydrazine (1.9
mL, 1.94 g, 60 mmol) in THF (500 mL) was heated under reflux for 4
h. The mixture was cooled, filtered and the solvent was evaporated
20 under reduced pressure. Saturated aqueous sodium hydrogen carbonate
was added and the mixture was extracted with ethyl acetate. The
combined organic fractions were washed with brine, dried (Na₂SO₄)
and the solvent was evaporated under reduced pressure. Acetic acid
(300 mL) was added and the mixture was heated under reflux for 15
25 min. The mixture was cooled and the solvent was evaporated under
reduced pressure. Saturated aqueous sodium hydrogen carbonate was
added and the mixture was extracted with ethyl acetate. The combined
organic fractions were washed with brine, dried (Na₂SO₄) and the
solvent was evaporated under reduced pressure to give 2,3-dihydro-5-
30 phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one as a foam (3.5 g, 52%).
δ_H (CDCl₃) 9.75 (1H, br s), 7.90-7.30 (5H, m), 6.87 (1H, d, J 6.0 Hz),
6.82 (1H, d, J 6.0 Hz), and 4.45 (2H, s).


- 72 -

Step C:10 2,3-Dihydro-1-methyl-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one

Sodium hydride (60% dispersion in mineral oil, 757 mg, 11.3 mmol) was added to a cooled (0°C) solution of 2,3-dihydro-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one (2.61 g, 10.8 mmol) in DMF (7 mL). Further DMF (10 mL) was added and the mixture was stirred for 30 min. A solution of iodomethane (0.67 mL, 1.53 g, 10.8 mmol) in ether (20 mL) was added and the mixture was stirred for 1 h. The mixture was poured into water and the mixture was extracted with ethyl acetate. The combined organic fractions were washed with brine, dried (Na₂SO₄) and evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with CH₂Cl₂/MeOH (95:5) to give 2,3-dihydro-1-methyl-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one (1.5 g, 54%).

δ H (CDCl₃) 7.67-7.35 (5H, m), 7.00 (1H, d, J 6.0 Hz), 6.85 (1H, d, J 6.0 Hz), 4.45 (2H, br s), and 3.50 (3H, s).

25

Step D:

30


- 73 -

3-Amino-2,3-dihydro-1-methyl-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one

2,3-Dihydro-1-methyl-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one (1.5 g, 5.8 mmol) was dissolved in toluene (30 mL).
5 The mixture was cooled to -10°C and potassium t-butoxide (1.7 g, 15.1 mmol) was added. The mixture was stirred at -10°C for 15 min., then isoamyl nitrite (1.0 mL, 0.87 g, 7.4 mmol) was added. The mixture was stirred at -10°C for 1 h. then allowed to warm to room temperature and poured into water (50 mL) and acetic acid (3 mL). The mixture
10 was extracted with ethyl acetate and the combined organic fractions were washed with brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with EtOAc/Hexane to give 2,3-dihydro-1-methyl-3-hydroxyimino-5-phenyl-1H-thieno[2,3-e]-
15 1,4-diazepin-2-one (0.80 g, 48%).

2,3-Dihydro-1-methyl-3-hydroxyimino-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one (0.80 g, 2.8 mmol) was dissolved in ethanol (40 mL) and Raney nickel (2 g) was added. The mixture was shaken under hydrogen (50 p.s.i.) for 5 days, adding further Raney
20 nickel (10 g) in portions. The mixture was filtered and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with CH₂Cl₂/MeOH to give 3-amino-2,3-dihydro-1-methyl-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one (248 mg, 33%).
25 ¹H (CDCl₃) 7.50-7.30 (5H, m), 7.05 (1H, d, J 6.0 Hz), 6.85 (1H, d, J 6.0 Hz), 4.57 (1H, s), 3.55 (3H, s), and 1.70 (2H, br s).

- 74 -

Step E:

15

1-(2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-3-yl)-3-(3-methylphenyl)urea

3-Methylphenylisocyanate (60 μ L, 62 mg, 0.46 mmol) was added to a solution of 3-amino-2,3-dihydro-1-methyl-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-2-one (124 mg, 0.46 mmol) in tetrahydrofuran (5 mL). The mixture was stirred at room temperature for 2 h. and the solvent was evaporated under reduced pressure. The residue was crystallized from EtOAc (4 mL) to give 1-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-thieno[2,3-e]-1,4-diazepin-3-yl)-3-(3-methylphenyl)urea as a solid (94 mg, 50%).

m.p. 128-130°C.

δ H (CDCl₃) 8.70 (1H, s), 7.65-6.75 (12H, m), 5.55 (1H, d, J 9.0 Hz), 3.55 (3H, s), and 2.30 (3H, s).


Anal. Calcd. for C₂₂H₂₀N₄O₂S.0.25H₂O:

C, 64.62; H, 4.99; N, 13.70.

Found: C, 64.68; H, 4.96; N, 13.70%.

30

- 75 -

EXAMPLE 55

5
10 (+)-N-[(3R)-7-Amino-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide

Step A:

15 To a mixture of 3(R)-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (*J. Org. Chem.* 1987, 52:3232-3239) (3.98 g, 15.0 mmol) in concentrated sulfuric acid (15 mL) cooled in an ice-bath was added dropwise a solution of potassium nitrate (2.12 g, 21.0 mmol) in concentrated sulfuric acid (6 mL). The mixture was stirred with cooling for 2 h., then stirred at ambient temperature for
20 1.5 h. Ice (80 g) was added and the mixture was basified with concentrated ammonium hydroxide to pH 9. The resulting mixture was extracted with ethyl acetate (3 x 220 mL). The combined organic fractions were washed with brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was purified by flash
25 column chromatography on silica gel, eluting with chloroform/methanol (97:3). The material which eluted was further purified by flash column chromatography on silica gel, eluting with ethyl acetate/methanol (95:5). The material which eluted was stirred under *n*-butyl chloride (30 mL) and the solvent was evaporated under reduced pressure to give
30 an inseparable mixture of 3(R)-amino-1,3-dihydro-1-methyl-7-nitro-5-phenyl-2H-1,4-benzodiazepin-2-one and 3(R)-amino-1,3-dihydro-1-methyl-7-nitro-5-(2-nitrophenyl)-2H-1,4-benzodiazepin-2-one (3.81 g) in a 3:1 ratio as a yellow solid.

- 76 -

δ H (CDCl₃) (mononitro compound) 8.43 (1H, dd, J 9, 3 Hz), 8.23 (1H, d, J 3 Hz), 7.59 (2H, m), 7.52 (2H, m), 7.44 (2H, m), 4.47 (1H, s), 3.53 (3H, s), and 2.42 (2H, br s); (dinitro compound) 8.49 (1H, dd, J 9, 3), 8.42 (1H, m), 8.18 (1H, d, J 3 Hz), 8.01 (1H, m), 7.67 (1H, t, J 6 Hz), 7.6-7.4 (2H, m), 4.52 (1H, s), 3.56 (3H, s), and 2.42 (2H, br s).

Step B:

A solution of 3-(2,4-dichlorophenyl)propionic acid (482 mg, 2.2 mmol), DMF (0.017 mL, 0.22 mmol), and thionyl chloride (0.24 mL, 3.3 mmol) in chloroform (2.5 mL) was heated at reflux for 1 h. The solvent was evaporated under reduced pressure to give 3-(2,4-dichlorophenyl)propionyl chloride (520 mg, 100%). To a solution of mixed 3(R)-amino-1,3-dihydro-1-methyl-7-nitro-5-phenyl-2H-1,4-benzodiazepin-2-one and 3(R)-amino-1,3-dihydro-1-methyl-7-nitro-5-(2-nitrophenyl)-2H-1,4-benzodiazepin-2-one (3:1) (621 mg, 2 mmol) and triethylamine (0.305 mL, 2.2 mmol) in methylene chloride (10 mL), was added a solution of 3-(2,4-dichlorophenyl)propionyl chloride (520 mg, 2.2 mmol) in methylene chloride (1.5 mL). The mixture was stirred for 30 min., the solvent was partially evaporated under reduced pressure, and the reaction mixture was purified by flash column chromatography on silica gel, eluting with methylene chloride/ether (90:10) to give a mixture of (+)-N-[(3R)-2,3-dihydro-1-methyl-7-nitro-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide and (+)-N-[(3R)-2,3-dihydro-1-methyl-7-nitro-2-oxo-5-(2-nitrophenyl)-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide (850 mg, 84%) in a 3:1 ratio as a solid white foam.

δ H (CDCl₃) (mononitro compound) 8.45 (1H, dd, J 9, 3 Hz), 8.25 (1H, d J 3 Hz), 7.54 (3H, m), 7.45 (2H, m), 7.38 (1H, d, J 2 Hz), 7.26-7.18 (4H, m), 5.50 (1H, d, J 8 Hz), 3.52 (3H, s), 3.10 (2H, m), and 2.70 (2H, m); (dinitro compound) 8.51 (1H, dd, J 9, 3 Hz), 8.40 (1H, m), 8.21 (1H, d J 3 Hz), 7.98 (1H, m), 7.68 (1H, t, J 6 Hz), 7.60 (1H, m), 7.44 (1H, m), 7.26-7.15 (4H, m), 5.52 (1H, d, J 8 Hz), 3.55 (3H, s), 3.10 (2H, m), and 2.70 (2H, m).

- 77 -

Step C:

To a solution of mixed N-[(3R)-2,3-dihydro-1-methyl-7-nitro-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide and (+)-N-[(3R)-2,3-dihydro-1-methyl-7-nitro-2-oxo-5-(2-nitrophenyl)-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide (3:1) (770 mg, 1.5 mmol) in acetic acid (6 mL) was added dropwise in portions over 1.5 h. a solution of 15% titanium (III) chloride in 20-30% hydrochloric acid (7.8 mL, 9.0 mmol). The resulting solution was stirred 30 min., basified with 20% sodium hydroxide solution (pH 9), diluted with water (80 mL) and extracted with ethyl acetate (3 x 100 mL). The combined organic fractions were washed with brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with ethyl acetate/hexane (75:25 increasing to 100:0). The first compound to elute was crystallized from ethyl acetate to give (+)-N-[(3R)-7-amino-2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide (413 mg, 57%) as a pale yellow solid, m.p. 179-180°C, [α]_D +60.2° (c= 0.500, CHCl₃).

δ H (CDCl₃) 7.60 (2H, d, J 7 Hz), 7.49-7.36 (5H, m) 7.24 (1H, d, J 9 Hz), 7.17 (2H, m), 6.99 (1H, dd, J 9, 3 Hz), 6.64 (1H, d, J 3 Hz), 5.54 (1H, d, J 8 Hz), 4.80-3.50 (2H, br s), 3.39 (3H, s), 3.09 (2H, t, J 8 Hz), and 2.68 (2H, dt, J_d 3, J_t 8 Hz).

Anal. Calcd. for C₂₅H₂₂Cl₂N₄O₂:

25 C, 62.38; H, 4.61; N, 11.64.

Found: C, 62.58; H, 4.68; N, 11.65%.

The second compound to elute was crystallized from ethyl acetate to give (+)-N-[(3R)-7-amino-2,3-dihydro-1-methyl-2-oxo-5-(2-aminophenyl)-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide (114 mg, 15%) as a pale yellow solid, m.p. 188-189°C, [α]_D +50.0° (c=0.100, MeOH).

δ H (CDCl₃) 7.36 (2H, m), 7.25 (1H, d, J 9 Hz), 7.15 (3H, m), 7.00 (1H, m), 6.88 (2H, m), 6.79 (1H, m), 6.60 (1H, bs), 5.52 (1H, d, J 8 Hz),

- 78 -


4.10-2.80 (4H br s), 3.40 (3H, m), 3.09 (2H, t, J 8 Hz), and 2.69 (2H, m).

Anal. Calcd. for $C_{25}H_{23}Cl_2N_5O_2 \cdot 0.05EtOAc$:

C, 60.43; H, 4.71; N, 13.99.

5 Found: C, 60.79; H, 4.74; N, 13.83%.

EXAMPLE 56



N-(2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-pyrido[4,3-e]-1,4-diazepin-3-yl)-3-(cyclohexyl)propanamide

A solution of dicyclohexylcarbodiimide (87 mg, 0.42 mmol) in methylene chloride (0.17 mL) was added to a solution of 3-amino-2,3-dihydro-1-methyl-5-phenyl-1H-pyrido[4,3-e]-1,4-diazepine-2-one (93 mg, 0.35 mmol) and cyclohexanepropionic acid (0.065 mL, 0.38 mmol) in tetrahydrofuran (0.5 mL) under argon. The resulting mixture was stirred for 5 h., filtered, and the filtrate was evaporated under reduced pressure. The residue was purified by preparative plate chromatography on silica gel eluting with methanol/chloroform/acetic acid (5:95:1). The purified material was stirred under chloroform (5 mL) with potassium carbonate (0.1 g) and water (2 drops) for 5 min. The mixture was dried (Na_2SO_4) and the solvent was evaporated under reduced pressure. The residue was crystallized from toluene to give N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-pyrido[4,3-e]-1,4-diazepin-3-yl)-3-(cyclohexyl)propanamide (47 mg, 33%) as a white crystalline solid, m.p. 170-173°C.

20

25

30

δ_H ($CDCl_3$) 8.75 (1H, d, J 6 Hz), 8.61 (1H, s), 7.58 (2H, m), 7.52 (1H, m), 7.45 (2H, m), 7.31 (1H, d, J 6 Hz), 7.21 (1H, d, J 8 Hz), 5.54 (1H,

- 80 -

solution and brine (20 mL each), dried (Na₂SO₄), and the solvent was evaporated under reduced pressure. The residue was triturated with hexane and the solid was collected and dried *in vacuo* to give 2,3-dihydro-3-hydroxyimino-1-methyl-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepine-2-one (1.60 g, 100%) as a tan foam.
5 δ H (CDCl₃) 8.77 (1H, s), 8.50 (1H, d, J 4 Hz), 7.81 (2H, dd, J 8, 1 Hz), 7.60 (1H, m), 7.49 (3H, m), 7.32 (1H, d, J 5 Hz), and 3.55 (3H, s).

Step B:

10 A solution of stannous chloride dihydrate (3.72 g, 16.5 mmol) in concentrated hydrochloric acid (11 mL) was added dropwise to 2,3-dihydro-3-hydroxyimino-1-methyl-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepine-2-one (1.54 g, 5.5 mmol) cooled in an ice bath. The resulting solution was stirred at ambient temperature for 3 h. The
15 solution was diluted with water (20mL), basified with concentrated ammonium hydroxide (18 mL), and extracted with ether (4 x 75 mL). The combined organic fractions were washed with brine (30 mL), dried (Na₂SO₄), and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica gel,
20 eluting with methanol/chloroform/acetic acid (5:95:1 increasing to 10:90:1). The material which eluted was stirred under chloroform (20 mL) with potassium carbonate (0.3 g) and water (2 drops) for 5 min. The mixture was dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was stirred under hexane, and the
25 resulting solid was collected to give 3-amino-2,3-dihydro-1-methyl-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepine-2-one (241 mg, 16%) as a yellow solid, m.p. 94-118°C.
 δ H (CDCl₃) 8.79 (1H, s), 8.48 (1H, d, J 5 Hz), 7.62 (2H, dd, J 8, 1 Hz), 7.51 (1H, m), 7.45 (2H, m), 7.24 (1H, dd, J 5, 1 Hz), 4.47 (1H, s), 3.55
30 (3H, s), and 2.2 (2H, bs).

Anal. Calcd. for C₁₅H₁₄N₄O.0.25(C₂H₅)₂O:

C, 67.46; H, 5.84; N, 19.67.

Found: C, 67.28; H, 5.66; N, 19.53%.

- 81 -

High res. mass spectrum: Theoretical mass for $C_{15}H_{14}N_4O$ (M+1): 267.124586. Measured mass (M+1): 267.123093.


Step C:

- 5 A solution of oxalyl chloride (0.023 mL, 0.26 mmol) in methylene chloride (0.2 mL) was added dropwise to a solution of 3-(2,4-dichlorophenyl)propionic acid (48 mg, 0.22 mmol) and DMF (1 drop) in methylene chloride (0.5 mL) cooled in an ice-bath. The resulting solution was stirred 1 h. with cooling. The solvent was
- 10 evaporated under reduced pressure to give 3-(2,4-dichlorophenyl)propionyl chloride (52 mg, 100%). To a solution of 3-amino-2,3-dihydro-1-methyl-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepine-2-one (53 mg, 0.20 mmol) and pyridine (0.021 mL, 0.22 mmol) in methylene chloride (3 mL), was added a solution of 3-(2,4-dichlorophenyl)propionyl chloride (52 mg, 0.22 mmol) in methylene chloride (0.5
- 15 mL). The mixture was stirred for 1 h., the solvent was partially evaporated under reduced pressure, and the reaction mixture was purified by flash column chromatography on silica gel, eluting with methanol/ether (5:95 increasing to 7.5:92.5). The material which eluted
- 20 was crystallized from toluene/hexane to give N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepin-3-yl)-3-(2,4-dichlorophenyl)propanamide (38 mg, 38%) as a white crystalline solid, m.p. 220-221°C.
- 25 δ_H (CDCl₃) 8.81 (1H, s), 8.52 (1H, d, J 5 Hz), 7.56 (2H, dd, J 7, 2 Hz), 7.51 (1H, m), 7.44 (2H, d, J 6 Hz), 7.40 (1H, m), 7.27 (2H, m), 7.18 (2H, dd, J 8, 2 Hz), 5.48 (1H, d, J 8 Hz), 3.55 (3H, s), 3.10 (2H, t, J 7 Hz), and 2.71 (2H, dt, J_d 2 J_t 8 Hz).
- Anal. Calcd. for $C_{24}H_{20}Cl_2N_4O_2 \cdot 0.25PhCH_3$:
C, 63.06; H, 4.52; N, 11.43.
- 30 Found: C, 63.03; H, 4.48; N, 11.25%.

- 82 -


EXAMPLE 58

N-[2,3-Dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide

Step A:

To a solution of the benzodiazepene (1.0 g, 5.3 mmol) in THF (20 mL) at -78°C under argon was added 60% NaH (0.252 g, 6.3 mmol) and the mixture stirred at -78°C for 1/2 hour. The reaction was then allowed to warm to 25°C and stirred for 2 hours before quenching into cold aq. NH₄Cl (10%) and extracting the product into ethyl acetate (3x50 mL). Concentration of the dried (Na₂SO₄) extracts gave an oil which was passed through silica (EtOAc/hexane) to give 1.35 g product (89%).

¹H NMR (CDCl₃) δ: 1.60 (s, 9H), 3.40 (s, 3H), 3.95 (brd, 1H), 4.80 (brd, 1H), 7.20 (d, 1H), 7.30 (q, 1H), 7.60 (t, 1H), 7.92 (d, 1H).

Step B:

- 79 -

d, J 8 Hz), 3.51 (3H, s), 2.39 (2H, m), 1.73 (4H, m), 1.63 (3H, m), 1.85-1.12 (4H, m), and 0.94 (2H, m).

Anal. Calcd. for C₂₄H₂₈N₄O₂·0.10PhCH₃:


C, 71.70; H, 7.02; N, 13.54.

5 Found: C, 71.78; H, 7.01; N, 13.57%.

Employing the procedure substantially as described above, but substituting 3-(4-trifluoromethylphenyl)-propionic acid for the cyclohexanepropionic acid, the following compound was prepared:

10

EXAMPLE 57



20 N-(2,3-Dihydro-1-methyl-2-oxo-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepin-3-yl)-3-(2,4-dichlorophenyl)propanamide

Step A:


25 To a solution of 2,3-dihydro-1-methyl-5-phenyl-1H-pyrido[3,4-e]-1,4-diazepine-2-one (*Can. J. Chem.* 1987, 65:1158-1161) (1.43 g, 5.7 mmol) in toluene (28 mL) under argon cooled to -20°C (ice/methanol bath) was added potassium t-butoxide (1.59 g, 14.2 mmol). The resulting purple suspension was stirred 15 min. at -20°C and isoamyl nitrite (0.92 mL, 6.8 mmol) was added. The mixture was
30 stirred at -20°C for 30 min., then poured into a mixture of water (25 mL), acetic acid (2.5 mL), and ethyl acetate (55 mL). The mixture was stirred to dissolve all solids and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 55 mL). The combined organic fractions were washed with saturated sodium bicarbonate

- 83 -

To a solution of the BOC-benzodiazepine (4.0 g, 13.8 mmol) in THF (80 mL) under argon was rapidly added a solution of isopropylmagnesium chloride (2.0 M) in THF (7.66 mL, 15.3 mmol). The reaction was stirred to 1/2 hour, quenched into aq NH₄Cl (50 mL), and extracted with ethyl acetate (2x200 mL). The organic extracts were concentrated and chromatographed on silica (1:1, EtOAc/hexane) to give 1.55 g (34%) of product.

¹H NMR (CDCl₃) δ: 1.14 (d, 3H), 1.19 (d, 3H), 1.40 (s, 9H), 3.13 (s, 3H), 3.2-3.8 (m, 3H), 5.45 (brs, 1H), 7.28 (dt, 1H), 7.48 (dt, 1H), 7.56 (dt, 1H), 7.72 (dd, 1H).

Step C:



To a 0°C solution of the isopropylphenone (1.55 g) in ethyl acetate was added anhydrous HCl gas over 90 min. The reaction was then concentrated in vacuo so a solid which was dissolved in H₂O (40 mL) and the pH adjusted to 11.0 with 1N LiOH. After 30 min. at pH = 11.0 the pH was adjusted to 7.0 with 1N HCl and product extracted into ethyl acetate. The organic extracts were dried (Na₂SO₄), filtered and concentrated to give a solid 1.22 g, 100%.

¹H NMR (CDCl₃) δ: 0.95 (d, 3H), 1.30 (d, 3H), 3.16 (septet, 1H), 3.36 (s, 3H), 3.60 (d, 1H), 4.60 (d, 1H), 7.2-7.3 (m, 2H), 7.45-7.55 (m, 2H).

Steps D-F:

The benzodiazepine was (D) converted to the corresponding oxime, (E) reduced and (F) coupled with (2,4-dichlorophenyl)-propionic acid as described in Example 80, Steps A, B

- 84 -

and C respectively to yield N-[2,3-dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)-propanamide.

m.p. 173-174°C

¹H NMR (CDCl₃) δ: 0.92 (d, 3H), 1.25 (d, 3H), 2.65 (dt, 2H), 3.05 (t, 2H), 3.15 (SepT, 1H), 3.40 (s, 3H), 5.38 (d, 1H), 7.0-7.6 (m, 8H).

The following compounds were prepared in a similar manner as described in Example 81.

10

EXAMPLE 59

N-[2,3-dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-cyclohexylpropanamide

15 m.p. 164-165°C

CHN: Anal. Calcd. for C, 71.51; H, 8.46; N, 11.37.

 Observed: C, 71.72; H, 8.39; N, 11.32.

EXAMPLE 60

20

N-[2,3-dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-(4-trifluoromethylphenyl)propanamide

m.p. 187-188°C

¹H NMR (CDCl₃) δ: 0.92 (d, 3H), 1.25 (d, 3H), 2.66 (dt, 2H), 3.04 (t, 2H), 3.15 (SepT, 1H), 3.40 (s, 3H), 5.38 (d, 1H), 7.14 (brd, 1H), 7.25-7.6 (m, 8H)

30

- 85 -

EXAMPLE 61

✓
N-[2,3-dihydro-1-methyl-2-oxo-5-(2-furanyl)-1H-1,4-benzo-
diazepin-3-yl]-3-cyclohexylpropanamide

5

m.p. 168-169°C

CHN: Anal. Calcd. for C, 70.21; H, 6.92; N, 10.68.
 Observed: C, 70.15; H, 6.67; N, 10.64.

EXAMPLE 62

✓ 10

N-[2,3-dihydro-1-methyl-2-oxo-5-(2-furanyl)-1H-1,4-benzo-
diazepin-3-yl]-3-(4-trifluoromethylphenyl)propanamide

15

m.p. 155-157°C

CHN: Anal. Calcd. for C, 63.29; H, 4.432; N, 9.23.
 Observed: C, 63.22; H, 4.44; N, 9.07.

EXAMPLE 63

20

N-[2,3-dihydro-1-methyl-2-oxo-5-(2-furanyl)-1H-1,4-benzo-
diazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide

m.p. 132-133°C

25


✓
CHN: Anal. Calcd. for C, 60.54; H, 4.20; N, 9.21.
 Found: C, 60.62; H, 4.07; N, 9.07.

EXAMPLE 64

30

N-[2,3,4,5-tetrahydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzo-
diazepin-3-yl]-3-cyclohexylpropanamide

- 86 -



A solution of N-[2,3-dihydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-cyclohexylpropanamide (50 mg) in methanol (10 mL), containing 10% Pd/C (50 mg) was stirred under 1 atmosphere of hydrogen for 18 hours. Filtration of the reaction, concentration and crystallization from diethyl ether gave 21 mg N-[2,3,4,5-tetrahydro-1-methyl-2-oxo-5-isopropyl-1H-1,4-benzodiazepin-3-yl]-3-cyclohexylpropanamide.


CHN: Anal. Calcd. for C, 71.12; H, 8.95; N, 11.31.
 Observed: C, 70.98; H, 8.97; N, 11.15.

m.p. 114-115°C

EXAMPLE 65


✓
30 N-[2,3-dihydro-1-methyl-2-oxo-5-methyl-1H-1,4-benzodiazepin-3-yl]-3-(2,4-dichlorophenyl)propanamide

- 87 -

Step A:

10 To CBZ-benzodiazapene (250 mg, 0.776 mmol) in toluene (25 mL) at reflux was added dropwise a solution of DMF dimethylacetal (1.09 mL) in toluene (10 mL). The reaction was refluxed for 5 hours, cooled and concentrated to an oil. The oil was triturated with ether to give a white solid (124 mg).


15 ¹H NMR (CDCl₃) δ: 2.50 (s, 3H), 3.42 (s, 3H), 5.12-5.20 (m, 3H), 6.62 (d, 1H), 7.25-6.4 (m, 7H), 7.5-7.6 (m, 2H).

Step B:

25 The CBZ-amine-N-methyl amide (190 mg) was treated with 30% HBr/AcOH (0.8 mL) for 1 hour at room temperature. The reaction mixture was poured into ether (10 mL) at 0°C and the solid filtered. Solid dissolved in 10% Aq. NaOH (5 mL) and CH₂Cl₂ (10 mL) and organic layer separated, dried (Na₂SO₄), filtered and concentrated to an oil (172 mg, 110%).

30 ¹H NMR (CDCl₃) δ: 2.42 (s, 3H), 3.05 (brs, 2H), 3.40 (s, 3H), 4.40 (s, 1H), 7.2-7.6 (m, 4H).


- 88 -

Step C:

20 N-[2,3-dihydro-1-methyl-2-oxo-5-methyl-1H-1,4-benzodiazepin-3-yl]-
3-(2,4-dichlorophenyl)propanamide was prepared in a similar manner
as described previously.
m.p. 194-195°C
CHN: Anal. Calcd. for C, 59.42; H, 4.74; N, 10.39.
 Observed: C, 59.50; H, 4.74; N, 10.44.
25 ¹H NMR (CDCl₃) δ: 2.49 (brs, 3H), 2.65 (dt, 2H), 3.05 (t, 2H), 3.42 (s,
3H), 5.35 (d, 1H), 7.1-7.6 (m, 8H).

30

- 89 -

EXAMPLE 66

5
10 3R-(+)-3-(Phenylthio)-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-propanamide

To a stirred solution of 3-Bromopropionic acid (1.0 g, 6.5 mmol) in DMF (20 mL) was added K₂CO₃ (1.8 g, 13 mmol) and thiophenol (0.72 g, 6.5 mmol). This was heated to 50°C for 1h. The
15 rm was then diluted with 200 mL H₂O and extracted with 2 x 100 mL EtOAc. The combined organics were washed with 100 mL H₂O and dried with Na₂SO₄. This was evaporated to 1.52 g of a colorless oil, 1.18 g corrected for residual DMF by NMR.


The above oil was taken up in 30 mL DMF and 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.45 g, 12.8
20 mmol) and 1-Hydroxybenzotriazole hydrate (1.73 g, 12.8 mmol) were added. This was stirred for 5 min at rt. 3-(R)-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (0.66 g, 2.6 mmol) was then added and the reaction was stirred at rt overnight. The reaction
25 was diluted with 200 mL H₂O and extracted with 2x150 mL EtOAc. The combined organics were washed with 1x100 mL H₂O, dried with Na₂SO₄ and evaporated. The residue was chromatographed over silica eluting with 2% MeOH:CHCl₃. Collected pure fractions, evaporated. Evaporated from diethyl ether to give 770 mg of a white foam.

30 Anal. Calcd for C₂₅H₂₃N₃O₂S•0.05Hexane:

C, 70.04; H, 5.51; N, 9.69.

Found: C, 69.91; H, 5.40; N, 9.78.

- 90 -

EXAMPLE 67

3R-(+)-5-(Methylthio)-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-propanamide

To an aqueous solution of K_2CO_3 (0.76 g, 5.5 mmol) was added 5-Bromopentanoic acid and sodium thiomethoxide. This was stirred at rt overnight. The reaction was diluted with 50 mL H_2O and acidified to pH=0 with 6N HCl. Extracted with 2 x 50 mL EtOAc. Dried with Na_2SO_4 , evaporated to 0.55 g of a yellow oil.


The above oil was taken up in 10 mL DMF and 1-(3-Dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride (1.30 g, 6.8mmol) and 1-Hydroxybenztriazole hydrate (0.92 g, 6.8 mmol) were added. 3-(R)-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (0.85 g, 3.4 mmol) was then added and the reaction was stirred overnight at rt. The reaction was diluted with 100 mL H_2O and extracted with 2 x 50 mL EtOAc. Combined organics were dried with brine and Na_2SO_4 , and evaporated to yellow oil. The residue was chromatographed over silica eluting with 50:50 EtOAc:Hex to 100% EtOAc. Pure fractions collected to give 1.33g of a colorless oil. 0.4 g of which were chromatographed over silica eluting with 2% MeOH: CH_2Cl_2 . Pure fractions were collected, evaporated. Evaporated from ethyl ether:hexane to give a white powder mp. 61-65°C.

Anal. Calcd for $C_{22}H_{25}N_3O_2S \cdot 0.35H_2O$:

C, 65.76; H, 6.45; N, 10.46.

Found: C, 65.81; H, 6.21; N, 10.57.

- 91 -

EXAMPLE 68

25 N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-N-2-(2-methoxyethoxy)-ethylhexanamide

3-(R)-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (1.33 g, 5.0 mmol) in N,N-dimethyl formamide (30 mL) was mixed with 1-bromo-2-(2-methoxyethoxy)-ethane (1.35 mL, 5.0 mmol) and triethylamine (1.0 mL). The mixture was stirred and heated at reflux for 4 h. After that time, the reaction was cooled to room temperature, diluted with 150 mL water, and extracted with ethyl acetate (3 x 100 mL). The organic layers were combined, dried with magnesium sulfate, gravity filtered, and the solvent was removed *in*

30

- 92 -

vacuo. The resulting oil was chromatographed over silica in 1:1 ethyl acetate:hexane, yielding the mono-alkylated product (1.2 g, 65%) as well as the starting 1,4-benzodiazapin-2-one and bis-alkylated material. To a solution of the mono-alkylated material (1.2 g, 3.27 mmol) in
5 methylene chloride (20 mL) was added hexanoyl chloride (0.96 mL, 3.27 mmol) and the reaction was stirred for 0.25 h. The reaction was then diluted with methylene chloride (150 mL) and saturated aqueous sodium hydrogen carbonate (150 mL) was added. The aqueous portion was extracted again with methylene chloride (2 x 100 mL) and the
10 organics were combined, dried with magnesium sulfate, gravity filtered, and the solvent was removed *in vacuo*. The resulting oil was chromatographed over silica with 1:1 ethyl acetate:hexane, yielding an oil, giving 580 mg (38%) of the product.
[α]_D 0.00°; c=0.27; MeOH.


15 Anal. Calcd. for C₂₇H₃₅N₃O₄•0.80 mol H₂O:
C, 67.56; H, 7.69; N, 8.75.
Found: C, 67.56; H, 7.39; N, 8.85%.

20

25

30

- 93 -

EXAMPLE 69

25 (+)-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-N-(5-hydroxypentyl)hexanamide

30 3-(R)-Amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one (1.33 g, 5.0 mmol) in acetonitrile (40 mL) was mixed with 5-chloropentan-1-ol (0.61 g, 5.0 mmol) and sodium hydrogen carbonate (2.0 g) was suspended in the mixture. The mixture was stirred and heated at reflux for 12 h. After that time, the reaction was cooled to room temperature, diluted with 100 mL water, and extracted with ethyl acetate (3 x 75 mL). The organic layers were combined, dried with magnesium sulfate, gravity filtered, and the solvent was removed *in vacuo*. The resulting oil was chromatographed

- 94 -


over silica in 1:49 methanol:chloroform yielding the mono-alkylated product (1.1 g, 62%) as well as the starting 1,4-benzodiazapin-2-one and bis-alkylated material. To a solution of the monoalkylated material (0.50 g, 1.42 mmol) in methylene chloride (30 mL) was added hexanoyl chloride (0.20 mL, 1.42 mmol) and the reaction was stirred for 0.25 h. The reaction was then diluted with methylene chloride (100 mL) and saturated aqueous sodium hydrogen carbonate (100 mL) was added. The aqueous portion was extracted again with methylene chloride (2x75 mL) and the organics were combined, dried with magnesium sulfate, gravity filtered, and the solvent was removed *in vacuo*. The resulting oil was chromatographed over silica with 1:1 ethyl acetate:hexane, yielding a foam, giving 360 mg (64%) of the product. foam, $[\alpha]_D^{25} (+)8.36^\circ$ ($c=0.61$, MeOH).

Anal. Calcd. for $C_{27}H_{35}N_3O_2 \cdot 0.25 \text{ mol H}_2O$:

Found: C, 71.42; H, 7.88; N, 9.25.

Found: C, 71.47; H, 7.89; N, 9.12%.

EXAMPLE 70



(+)-3-Cyclohexyl-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-N-(tetrazolylmethyl)propanamide

(+)-N-[2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-N-(hydroxymethyl)hexanamide (0.67 g, 1.56 mmol) was dissolved in methylene chloride (100 mL), along with tetrazole (0.33 g, 4.7 mmol), and then, N,N-diisopropyl-dibenzylphosphoramidite (1.07 g, 3.1 mmol). After 2 h, the mixture was diluted with methylene chloride (150 mL), and extracted with saturated

- 95 -

aqueous sodium hydrogen carbonate (3 x 100 mL). The organic layers were combined, dried with magnesium sulfate, gravity filtered, and the solvent was removed *in vacuo*. The resulting oil was chromatographed twice over silica with 1:1 ethyl acetate:hexane, yielding two constitutional isomers, a (65 mg, 9%) and b (56 mg, 7.5%).

Isomer A:

m.p. 96-98°C, $[\alpha]_D +188.9^\circ$ (c=0.19, MeOH).Anal. Calcd. for $C_{27}H_{31}N_7O_{20.30}$ mol TFA:

10 C, 63.78; H, 6.07; N, 18.86.

Found: C, 63.7; H, 6.12; N, 18.76%.

Isomer B:

m.p. 92-95°C, $[\alpha]_D +81.3^\circ$ (c=0.31, MeOH).Anal. Calcd. for $C_{27}H_{31}N_7O_{20.35}$ mol TFA:

15 C, 63.31; H, 6.01; N, 18.66.

Found: C, 63.35; H, 6.02; N, 18.74%.

20

25

30

- 96 -

WHAT IS CLAIMED IS:

1. A method of treating cardiac arrhythmia in mammals comprising block of the slowly activating delayed rectifier potassium (K⁺) current (I_{Ks}) and the rapidly activating and deactivating delayed rectifier potassium current (I_{Kr}).
5
2. The method of Claim 1 wherein the slowly activating delayed rectifier potassium (K⁺) current (I_{Ks}) in isolated myocytes is selectively blocked by a compound at a concentration of 1 μM or less (IC₅₀) and the concentration that blocks I_{Ks} by 50% is at least 10 fold lower than the concentration required to cause 50% block of I_{Kr} and/or I_{K1}.
10
3. The method of Claim 2 wherein the block prolongs the cardiac refractory period equally or to a greater extent as heart rate increases.
15
4. The method of Claim 3 wherein the block is independent of heart frequency.
20
5. The method of Claim 1 wherein the blockade is induced through the use of a 1,4-benzodiazepine or benzodiazepine derivative.
25
6. The method of Claim 5, wherein the benzodiazepine or benzodiazepine derivatives selectively block the slowly activating delayed rectifier potassium (K⁺) current (I_{Ks}).
30
7. The method of Claim 6, wherein the 1,4-benzodiazepines or benzodiazepine derivatives are selected from the group consisting of those compounds which block the I_{Ks} current measured in isolated myocytes by 50% at a concentration of 1 μM or less

- 97 -

and exhibit a selectivity ratio of greater than 10 over blockade of IKr, IK1 and ICa.

8. The benzodiazepine or benzodiazepine derivative of Claim 5, selected from the group consisting of 3(R)-(+)-1,3-dihydro-3-(2-indolecarbonylamino)-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one; (S)-(-)-1,3-dihydro-3-(2-indolecarbonylamino)-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one; N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea; (R)-N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea; N-[3(R,S)-5-cyclohexyl-2,3-dihydro-1-methyl-2-oxo-1H-1,4-benzodiazepin-3-yl]-N'-[3-(isopropylsulfonylamino-carbonyl)phenyl]urea; N,N-dimethyl-4-(3(R,S)-(((3-methylphenyl)amino)-carbonyl)amino-1,3-dihydro-1-(2-methylpropyl)-2-oxo-1,4-benzodiazepin-5-yl)phenylmethanamine; (R)-3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one; N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea; (+-)-N-(2,3-dihydro-1-methyl-2-oxo-5-(4-pyridinyl)-1H-1,4-benzodiazepin-3-yl)-1H-indole-2-carboxamide; N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-4-methylbenzenesulfonamide; (R)-1-(2-(dimethylaminoethyl)-5-(2-fluorophenyl)-1,3-dihydro-3-((1-methyl-1H-indole-3-yl)methyl)-2H-benzodiazepin-2-one; N-(2,3-dihydro-1-methyl-2-oxo-5-(4-morpholino)-1H-1,4-benzodiazepin-3-yl)-N'-3-methylphenylurea.

25

9. A composition useful in the treatment of cardiac arrhythmia in mammals comprising a pharmaceutical compound and a suitable pharmaceutical carrier wherein the compound provides a selective block of the slowly activating delayed rectifier potassium (K+) current (IKs).

30

10. The composition of Claim 9, wherein the pharmaceutical compound is a 1,4-benzodiazepine or benzodiazepine derivative.

- 98 -

11. The composition of Claim 10, wherein the 1,4-benzodiazepine or benzodiazepine derivative is selected from the group consisting of 3(R)-(+)-1,3-dihydro-3-(2-indolecarbonylamino)-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one; (S)-(-)-1,3-dihydro-3-(2-indolecarbonylamino)-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one; N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea; (R)-N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(3-methylphenyl)urea; N-[3(R,S)-5-cyclohexyl-2,3-dihydro-1-methyl-2-oxo-1H-1,4-benzodiazepin-3-yl]-N'-[3-(isopropylsulfonylaminocarbonyl)-phenyl]urea; N,N-dimethyl-4-(3(R,S)-(((3-methylphenyl)amino)-carbonyl)amino-1,3-dihydro-1-(2-methylpropyl)-2-oxo-1,4-benzodiazepin-5-yl)phenylmethanamine; (R)-3-amino-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one; N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-N'-(m-methylphenyl)urea; (+-)-N-(2,3-dihydro-1-methyl-2-oxo-5-(4-pyridinyl)-1H-1,4-benzodiazepin-3-yl)-1H-indole-2-carboxamide; N-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl)-4-methylbenzenesulfonamide; (R)-1-(2-(dimethylaminoethyl)-5-(2-fluorophenyl)-1,3-dihydro-3-((1-methyl-1H-indole-3-yl)methyl)-2H-benzodiazepin-2-one; N-(2,3-dihydro-1-methyl-2-oxo-5-(4-morpholino)-1H-1,4-benzodiazepin-3-yl)-N'-3-methylphenylurea.

25

30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/13364

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/55

US CL :514/221, 821

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/221, 821

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS AND CAS ONLINE: compounds of the claims with arrhythmia, potassium current

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,820,834 (EVANS ET AL.) 11 APRIL 1989, see entire document.	9-11
Y	EP, A, 0,434,360 (Bock et al) 26 JUNE 1991, see entire document.	9-11
Y	EP, A, 0,434,369 (Bock et al) 26 JUNE 1991, see entire document.	9-11



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

&

document member of the same patent family

Date of the actual completion of the international search

12 JANUARY 1995

Date of mailing of the international search report

07 FEB 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KIMBERLY R. JORDAN

Telephone No. (703) 305-1235